

Yield and yield components of tomato grafted plants in the high Andean region of Colombia

Rendimiento y componentes del rendimiento en plantas injertadas de tomate en la región alto Andina en Colombia

JAMER ALEXIS RAMÍREZ-JIMÉNEZ^{1, 2}

CARLOS FELIPE BARRERA-SÁNCHEZ¹

OSCAR DE JESÚS CÓRDOBA-GAONA¹

¹ Universidad Nacional de Colombia, Facultad de Ciencias Agrarias, Medellín (Colombia). ORCID Ramírez-Jiménez, J.A.: <https://orcid.org/0000-0001-7768-1631>; ORCID Barrera-Sánchez, C. F.: <https://orcid.org/0000-0002-5015-2956>; ORCID Córdoba-Gaona, O.J.: <https://orcid.org/0000-0003-2642-4146>

² Corresponding author. jaaramirezji@unal.edu.co



Tomato grafted plant.
Photo: O.J. Córdoba-Gaona

Last name: RAMÍREZ-JIMÉNEZ / BARRERA-SÁNCHEZ / CÓRDOBA-GAONA

Short title: YIELD OF TOMATO GRAFTED PLANTS

Doi: <https://doi.org/10.17584/rcch.2020v14i3.11671>

Received for publication: 01-09-2020 Accepted for publication: 19-10-2020

ABSTRACT

Grafting as a technique is gaining full attention, and the production and demand for grafted vegetable plants have increased across the whole world, especially for greenhouse cultivation. The study aims to evaluate the potential of different scion x rootstock combinations or improving the yield of the tomato crop under a plastic house condition. For this purpose, a scion of tomato cv. Libertador was grafted on two commercial tomato rootstock ('Olimpo' and 'Armada') and cultivated in a plastic house in a randomized complete block experimental design, with four scion-rootstock combinations: vigor and resistance rootstocks, self-grafting and non-grafted plants. Yield and yield components were evaluated (number and weight of the fruits of extra, commercial, and non-commercial quality; so, weight per cluster, and weight per fruit). It was shown that, although vigor rootstocks produce less fresh fruit in the first harvests, from the seventh harvest onwards, the vigor rootstocks outperformed the other treatments in the accumulated yield by producing 37, 22 and 22% more yield fruit, and 60, 30 and 40% more number of fruits of extra quality, regarding resistance rootstock, self-grafted, and non-grafted plants. The fruits of vigor rootstock, self-grafted and non-grafted plants above 150 g tend to have a cylindrical shape; however, those of resistance rootstock tend to be flattened. The use of a vigor rootstock increased the yield of cv. Libertador, regarding the rootstock with resistance characteristics and controls, self-grafted, and non-grafted plants.

Additional key words: *Solanum lycopersicum*; scion-rootstock interaction; grafting; vigor; resistance.

RESUMEN

La injertación es una técnica que ha ido incrementando en todo el mundo, especialmente en el cultivo de tomate en invernadero. El objetivo de este trabajo fue evaluar los componentes de rendimiento y el rendimiento de un cultivar comercial de tomate injertado sobre diferentes combinaciones de patrones de la región altoandina de Colombia; para ello el cultivar Libertador fue injertado sobre los patrones 'Olimpo' y 'Armada', y como controles se utilizaron plantas auto injertadas y no injertadas del mismo cultivar. Se empleó un diseño de bloques al azar con cuatro

repeticiones y cuatro tratamientos: patrón vigor ('Olimpo'), patrón resistencia ('Armada'), autoinjerto y no injertadas. Hubo diferencias significativas entre tratamientos para el rendimiento acumulado. Las plantas de tomate injertadas en el portainjerto vigor produjeron 37, 22 y 22% más de peso de fruto, y 60, 30 y 40% más de frutos de calidad extra, en comparación con los tratamientos patrón de resistencia, plantas auto injertadas y no injertadas. Se observó que, si bien el uso de un patrón de vigor condujo a una menor producción de fruta en las primeras cosechas, a partir de la séptima cosecha, este tratamiento fue superior a los demás, en cuanto al rendimiento acumulado. Los frutos en el tratamiento vigor, auto injertos y plantas no injertadas que superaron los 150 g tienden a tener forma cilíndrica; mientras que, los frutos del tratamiento resistencia tienden a presentar una forma achatada. El uso de un portainjerto vigoroso aumentó el rendimiento del cv. Libertador, con respecto a los tratamientos portainjerto con características de resistencia y a los tratamientos control, plantas auto injertadas y no injertadas.

Palabras clave adicionales: *Solanum lycopersicum*; interacción portainjerto-patrón; injertación; vigor; resistencia.

INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is one of the most important and popular vegetable crops in the world, which has generated a large number of studies, focused on increasing the quality and yield of fruit and reducing the effects of different types of stress (Geboloğlu *et al.*, 2011; Singh *et al.*, 2017; Milenković *et al.*, 2020). Tomato cultivation can be carried out both in the open field and in protected conditions, where there are both biotic and abiotic limitations that affect growth, development, and fruit production (Djidonou *et al.*, 2016; Singh *et al.*, 2017). As one of the strategies to respond to the demands of the private and public sector, in order to overcome these types of limitations, grafting as a technique is gaining full attention, and the production and demand of grafted vegetable plants have increased across the whole world throughout the world, especially for greenhouse cultivation of vegetable crops (Geboloğlu *et al.*, 2011; Reddy, 2016; Kumar and Sanket, 2017). The grafting process involves the union of two parts (a rootstock and scion) together from two different plant parts to form a single living plant exhibiting the characters of both separately (Gaion *et al.*, 2018; Sen *et al.*, 2018). Initially, grafting was used widely in vegetable crops to limit the effects of soilborne pathogens; however, the utility of grafting has widened the

scope over the years. Grafted plants have been successfully used to induce tolerance against salinity (Singh *et al.*, 2020), drought (Fullana-Pericàs *et al.*, 2018), flooding (Bhatt *et al.*, 2015), temperature stress (Muneer *et al.*, 2016), and adverse chemical soil condition (Martínez-Andújar *et al.*, 2017), among other purposes. The new and improved root system provides tolerance through grafting, by deep and vigorous root formation (Aloni *et al.*, 2010), improved soil water and nutrient uptake (Savvas *et al.*, 2017) and increasing plant vigor and extending the duration of economic harvest time (Reddy, 2016; Kumar *et al.*, 2017; Kumar and Sanket, 2017).

On the other hand, the effect of grafting on the yield of tomato plants is unclear, and some studies affirm that it increases (Pogonyi *et al.*, 2005; Khah *et al.*, 2006; Djidonou *et al.*, 2016; Milenković *et al.*, 2020), others that have shown a performance equal to their controls, and even unfavorable effects on yields have been found with certain combinations (Miskovic *et al.*, 2016; Milenković *et al.*, 2020). In each production process, there are components that condition tomato yield; the consideration and quantification of these components allow to estimate the yield and make decisions that guarantee higher production (Torres, 2017). In general, the differences in the yield when tomato grafted plants are used, are based on the increase in the size and quality of the fruits, without an increase in the number of fruits; likewise, it is related to the distribution of the harvest (Pogonyi *et al.*, 2005; Khah *et al.*, 2006; Djidonou *et al.*, 2016; Milenković *et al.*, 2020). Although grafting has become a useful tool to enhance many traits in horticultural crops, its role in improving the tomato crop performance in the High-Andean region is yet proving. We hypothesized that the use of grafted plants improves the tomato fruit yield, promoting the use of this technique for the Colombian tomato producers. The objective of the present study was to evaluate the potential of different scion x rootstock combinations or improving the yield of the tomato crop under a plastic house condition.

MATERIALS AND METHODS

This experiment was carried out under plastic house conditions, at the El Socorro farm, El Santuario, Antioquia, Colombia (6°6'55.8" N and 75°13'10.15" W, an elevation of 2.251 m), which is localized in the high-Andean region. Properties of the soil used in the experiment were sandy loam-clay-sandy textural class, pH (5.0), EC (0.06 dS/m), OC (5.8 %), phosphorus (66 mg kg⁻¹ soil), sulfur (53.2 mg kg⁻¹ soil), Ca (10.6 cmolc kg⁻¹), Mg (3.0 cmolc kg⁻¹), K (2.47 cmolc kg⁻¹),

ECEC ($16.5 \text{ cmolc kg}^{-1}$), Fe (74 mg kg^{-1}), Mn (9 mg kg^{-1}), Cu (9 mg kg^{-1}), Zn (5 mg kg^{-1}) and B (0.2 mg kg^{-1}). Based on previous related soil condition, tomato plant nutrition was applied by using the chemical fertilization: N (35.3 g), P_2O_5 (26 g), K_2O (57.3 g), MgO (22.3 g), CaO (33 g), S (14 g), B (0.4), Cu (0.03 g), Fe (0.04 g), Mn (0.01 g), Mo (0.001 g), Zn (0.65 g), and Si (15 g) per plant, applied through fertigation and distributed at planting, and 45, 80, and 100 d after planting.

The average air temperature was 19.8°C , and the absolute minimum and maximum temperatures were 10.1 and 43.4°C ; on the other hand, the average relative humidity was 77% , with a maximum and minimum of 89 and 65% . The maximum air temperature was above the maximum threshold (35°C) in the middle of the days of the experimental period, while the minimum temperature was always above the base temperature for the tomato crop (10°C) (Fig. 1). The volume of water and frequency of tomato irrigation are shown in figure 2. The irrigation of the crop was carried out through drip irrigation, applying a total volume per plant of 243.2 L , and the average daily amount of water per plant was 1.42 L .




Figure 1. Daily mean temperature (Mean), maximum (MAX), minimum (MIN), and relative humidity (RH) into plastic house between 04/23/2019 to 10/25/2019.




Figure 2. Frequency of drip irrigation, and daily volume and accumulated volume of water applied by tomato plant between 04/23/2019 to 10/11/2019.

A randomized complete block experimental design was used, with four (4) treatments and four (4) repetitions. The treatments consisted of a single commercial tomato scion grafted on different rootstocks combination: vigor rootstock, resistance rootstock, self-grafting, and non-grafted plants. Each experimental plot consisted of 32 tomato plants, spread over four rows of eight plants each, for an area of 15.84 m².

The genotype (*S. lycopersicum* L.) used as a scion was a tomato Chonto cultivar Libertador; and as a rootstock, two commercial materials were used: ‘Olimpo’ as vigorous rootstock, and ‘Armada’ as disease resistant rootstock. The grafting methods applied were the tongue approach grafting, as described Lee *et al.* (2010) that consists in removed the growing point of the rootstocks and cut the rootstock in a downward direction and the scion cut in an upward direction at an angle perpendicular axis, and deep enough to allow the fusion of tongue insertion. After the graft is completed, specially designed clips are placed to fix the graft position. The non-grafted and grafted plants were transplanted to the plastic house on April 29, 2019, each of them, with the third leaf on the main shoot folded, corresponding to 103 stage according to the BBCH scale proposed by Feller *et al.* (1997). The field planting distance was 1.1 m between rows and 0.45 m between plants, and as part of the management, the first lateral shoot was allowed to grow below the first inflorescence in order to have two stems per plant, for a density of 20,200 plants/ha and 40,400 stems/ha. The

growth of the tomato plants was allowed until the ninth fruit cluster's emission on the main stem and seventh fruit cluster on the lateral stem, for a total of 16 fruit clusters emitted throughout the life cycle.

The harvest was carried out between 07/21/2019 to 10/28/2019, in total twelve harvests were made until the emission of the ninth cluster (C9) of fruits on the main stem (1) and the seventh cluster (C7) of fruits in the lateral stem (2), for a total of 16 fruit clusters emitted in the two stems (1C1, 1C2, 2C3, 1C4..., 2C15, 1C16) throughout the whole tomato life cycle. In each cluster, the number of fruits was determined for each cluster, and for each fruit, the parameters weight (g), equatorial diameter and longitudinal diameter were determinate. Finally, the number of fruits and fruit yield per cluster for the extra (>80 g), commercial (50 - 80 g) and non-commercial (<50 g) qualities, and total accumulated yield were estimated as the total accumulated weight of fruits.

One-way analysis of variance (ANOVA) was performed to reveal differences within each treatment (scion/rootstock combinations) for all measured parameters. For the mean comparison, an Honestly Significant Difference-HSD Test was applied, using the R project “agricolae” package (R Core Team, 2017). For the number of fruits and yield per cluster, a combined analysis by scion x rootstock combinations was performed. The variables longitudinal and equatorial diameter of tomato fruits and their relationship with weight per fruit were analyzed after a regression analysis using the Statgraphics program (5.1), for this, the dependent variables were DL and SD (DV) and two independent variables weight and treatment (VI). The procedure consisted of obtaining the least-quadratic equation that best expresses the relationship between the DV and VI and estimates the quality of the regression equation obtained using the coefficient of determination.

RESULTS AND DISCUSSION

The weight and number of fruits for each order of cluster harvested are presented in figure 3. The weight per cluster ($P>0.05$) and the number of fruits for each cluster ($P>0.05$) were not significantly affected by the scion x rootstock combinations, while the order of emission of the cluster significantly affected the weight and number of fruits. On the other hand, the yield and the number of fruits among the 16 clusters, independent of the grafting treatments, it was observed that the weight and number of fruits decrease with the order of emission, being the first clusters (1C1 and 1C2) of higher weight and number of fruits, values that decrease until the eighth cluster (1C8)

and from it up to the cluster 1C16, the clusters generally present similar values in the number and weight of the fruits (Fig. 3).




Figure 3. Weight and number of fruits for each tomato cluster for the different scion × rootstock combinations. Vigor rootstock (VR), resistance rootstock (RR), self-grafting (SELF), and non-grafted plants (NG). Values are mean ± standard error. *Significant difference at the 5 % level according to Honestly Significant Difference (HSD).

Pogonyi *et al.* (2005) found that grafting improved yields per cluster (eight clusters) when compared to non-grafted plants, this due to the production of a higher quantity of fruits and greater weight; and that the first clusters produce more significant quantities than the last ones, both in grafted and non-grafted plants.

Regarding the total number of fruits for each tomato quality, significant differences were found between the different scion x rootstock combinations for the extra quality ($P=0.0216$). In contrast,

for the commercial qualities ($P=0.388$) and non-commercial ($P=0.989$), there were no significant differences. The use of a vigor rootstock increased 38% more fruit of extra quality compared to the resistance rootstock and 28% more about self-grafted and non-grafted plants; on the other hand, the resistance rootstock, self-grafted, and non-grafted plants did not differ from each other for the number of extra quality fruits (Tab. 1). On the other hand, the number of tomato fruits for the extra quality represented 74% (vigor), 56% (resistance), 64% (self-grafted), and 62% (non-grafted plants) of the total fruits harvested in each treatment. On average, fruits of commercial quality represented 14% of the total fruits, while non-commercial quality was 19% for all treatments (Tab. 1). Regarding the weight per fruit, no differences were observed in each of the fruit qualities; nor in the total number of fruits due to different scion \times rootstock combinations (Tab. 1).

Table 1. The total number and weight of tomato fruits per qualities under different scion \times rootstock combinations.

Treatment	Fruit number per plant			Total
	NC	C	E	
VR	9.6 \pm 1.82 a	16.4 \pm 1.87 a	72.5 \pm 4.57 a	98.54 \pm 4.57 a
RR	19.5 \pm 5.07 a	21.4 \pm 4.18 a	52.9 \pm 4.14 b	93.8 \pm 6.10 a
SELF	15.5 \pm 2.37 a	17.5 \pm 1.76 a	58.9 \pm 0.92 b	91.3 \pm 3.14 a
NG	12.3 \pm 0.93 a	22.1 \pm 0.69 a	56.5 \pm 1.64 b	90.9 \pm 2.93 a
Treatment	Fruit weight by quality (g)			Mean fruit weight (g)
	NC	C	E	
VR	39.3 \pm 1.60 a	66.4 \pm 0.94 a	125.7 \pm 0.92 a	104.9 \pm 3.69 a
RR	38.5 \pm 0.94 a	65.4 \pm 1.41 a	118.9 \pm 3.04 a	89.3 \pm 4.85 a
SELF	37.8 \pm 0.17 a	64.4 \pm 0.61 a	121.9 \pm 4.97 a	97.2 \pm 3.95 a
NG	39.4 \pm 1.09 a	68.3 \pm 3.15 a	121.4 \pm 1.00 a	96.0 \pm 2.76 a

Extra (E), commercial (C), non-commercial (NC), vigor rootstock (VR), resistance rootstock (RR), self-grafting (SELF), and non-grafted plants (NG). Values are mean \pm standard error. Treatments with different low case letters are different at the 5 % level according to Honestly Significant Difference (HSD).

For the accumulated yield the resistance rootstock and the self-grafted and non-grafted plants produced significantly more fresh fruit (first five harvests) than rootstock vigor, however, from the eighth harvest onwards, rootstock vigor was highlighted significantly ($P=0.00143$) exceeding by 21 % the self-grafted and non-grafted plants treatments, and 36 % more than resistance rootstock (Fig. 4). Other authors observed similar results; the tomato yield in greenhouse conditions was higher in the self-grafted and non-grafted plants; however, in the end, the use of rootstocks of resistance and vigor significantly exceeded the yields (Pogonyi *et al.*, 2005; Khah *et al.*, 2006).

Which, according to Khah *et al.* (2006) is since grafted plants present stress due to the union of the parts (scion-rootstocks), which delays the maturity of the fruits and their harvest.




Figure 4. Accumulated tomato fruit yield for the different scion × rootstock combinations. Vigor rootstock (VR), resistance rootstock (RR), self-grafting (SELF), and non-grafted plants (NG). Harvest was performed between 07/21/2019 to 10/28/2019. Values are mean ± standard error. Treatments with different low case letters are different at the 5% level according to Honestly Significant Difference (HSD).

Tomato yield varied significantly between treatment for the extra ($P=0.0000626$), commercial ($P=0.00011$) and non-commercial qualities ($P=0.000000948$); likewise, the total yield per plant and area ($P=0.00118$) was significantly affected by the different grafting combinations. Regarding the contribution to total production, the extra (E) quality fruits represented 86% (9.7 kg/plant) of the yield of plants grafted on rootstock vigor, which was statistically different from 75% (6.01 kg/plant), 81% (7.4 kg/plant) and 77% (7.1 kg/plant) of the contribution in resistance rootstock, self-grafted, and non-grafted plants treatments respectively. The highest weight of commercial-grade (C) fruits was observed in the non-grafted plant treatment (1.5 kg/plant), while the highest yield in non-commercial (NC) quality was observed in the resistance rootstock combination (0.73 kg/plant) (Fig. 5).




Figure 5. Tomato fruit yield for the scion × rootstock combinations. Noncommercial (NC), Commercial (C), and Extra (E) fruit quality. Vigor rootstock (VR), resistance rootstock (RR), self-grafting (SELF), and non-grafted plants (NG). The harvest was performed between 07/21/2019 to 10/28/2019. Treatments with different low case letters are different at the 5% level according to Honestly Significant Difference (HSD).

Regarding the total yield by area, it was determined that when the scion was grafted on a vigorous rootstock, the yield was 40% higher compared to when it was grafted on a disease-resistant rootstock and 20% higher compared to both control treatments (self-grafted and non-grafted plants) (Fig. 5). The difference between the yield for each treatment vigor (43.9 kg m⁻²), resistance (32.0 kg) self-grafted (36.0 kg) and non-grafted (35.9 kg) plants is explained due to the more significant number of fruits and quality yield extra produced in the vigor treatment, which was of the order of 60, 30 and 40% more to resistance rootstock, self-grafted, and non-grafted plant treatments respectively (Fig. 5).

The higher yield of tomato plants depends on the proper selection of both the scion and the rootstock (Ntatsi *et al.*, 2014; Hossain *et al.*, 2019), in several studies results presented are not

precise; some indicate that the use of grafted plants increases yield; however, other authors report an adverse effect of the grafting on the performance of the commercial scion.

Adverse effects on performance were reported by Hossain *et al.* (2019) when comparing grafted plants with non-grafted and self-grafted controls; grafting reduced yields by 10% by area and 15% by the plant. Miskovic *et al.* (2016) when grafting tomato cultivar 'Jeremy' on eggplant rootstock cv. Madona observed an adverse effect of the eggplant rootstock by reducing the yield of cv. Jeremy, by 43%. Ntatsi *et al.* (2014), when grafting the commercial cv. Kommeet on the rootstock LA 1777, a significant reduction in yield was observed concerning the non-grafted and self-grafted controls. Finally, Riga (2015) evaluated the grafting of a commercial stem on nine different rootstocks and their respective controls (non-grafted and self-grafted) and although in the yield and number of fruits per area, and the average weight of the fruits they did not show statistical differences; Most of the rootstocks tended to be detrimental to the quality parameters considered.

Regarding positive effects on tomato yield in different scion x rootstock combinations, different results have been reported. 'Yanki F1' and 'Esin F1' scions grafted onto 11 rootstock combinations shown an increase in yield compared with self and non-grafted plants of up to 32.73% (Geboloğlu *et al.*, 2011). Qaryouti *et al.* (2007) also found that the fruit yield increased between 12 and 38% when grafting the cv. Cecilia on the 'He-Man' and 'Spirit' rootstocks. Djidonou *et al.* (2016) found a positive effect on the performance of the cultivar 'Florida 47' when it was grafted on the 'Beaufort' or 'Multifort' rootstocks compared to the non-grafted and self-grafted controls. Hossain *et al.* (2019), when comparing grafted plants with the self-grafted and non-grafted controls, found that the use of a rootstock improved the yield by area by 50% and by the plant by 56%. According to Rahmatian *et al.* (2014), the yield per plant increased significantly by 27%, in the grafted plants compared to their self-grafted and non-grafted plants.

The increase in the yield of the grafted plants is due to several factors among them: a higher quantity of fruit of extra and commercial quality (Qaryouti, *et al.*, 2007; Geboloğlu *et al.*, 2011), an increase in the total number of fruits (Turhan *et al.*, 2011), in the number of fruits per cluster (Ntatsi *et al.*, 2014), in the weight of fruits (Rahmatian *et al.*, 2014; Djidonou *et al.*, 2016) and the yield per plant (Rahmatian *et al.*, 2014). In addition to the above, the increase in yield in grafted plants is also attributed to a higher functional relationship between the rootstock and the stem in terms of hormones, proteins and other metabolites (Aloni *et al.*, 2010; Ntatsi *et al.*, 2014); as well

as higher capacity for absorbing water and minerals due to a more vigorous root system due to the use of the rootstock (Pogonyi *et al.*, 2005).

The linear regression model was significant ($P=0.0000$) for the fruit's weight as a function of the equatorial and longitudinal diameter, with a significant regression coefficient for the two diameters in all treatments. Figure 6 shows the linear regressions of fruit weight as an independent variable, and the respective equatorial diameters (ED) and longitudinal diameters (LD) as weight-dependent variables.




Figure 6. Linear regression models for equatorial diameter (ED) and longitudinal diameter (LD) depending on fruit weight, among the different scion \times rootstock combinations. Vigor rootstock (VR), resistance rootstock (RR), self-grafting (SELF), and non-grafted plants (NG).

Figure 6 shows that the equatorial and longitudinal diameter reach an approximate value of 6 cm when the fruits have a weight close to 150 g, indicating that tomato fruits have a circular or round shape. However, an increase in the longitudinal regarding equatorial diameter is observed for the resistance rootstock treatment at values higher than 150 g, which indicates that the fruits that exceed this weight tend to have a flatter shape compared to other treatments. In the vigor rootstock, self-grafted, and non-grafted plants, conversely, the increase in both diameters is similar, indicating that the fruits tend to have a round shape. When the Yeni Talya, Swanson, and Beril cultivars were grafted on the Beaufort and Arnold standards, the fruits became more flattened with any combination of grafting compared to the non-grafted control (Turhan *et al.*, 2011).

CONCLUSION

The vigorous rootstock increases the tomato yield of the cv. Libertador due to the higher yield and number of extra quality fruits, regarding the controls self-grafted and non-grafted plants. The scion x resistance rootstock combination affected tomato yield compared to self-grafted plants. The fruits of cv. Libertador grafted on resistance patterns presented flattened shape after 150 g of weight, while the shape of the fruit in plants grafted on vigor rootstock, self-grafted and non-grafted plants presented a round shape.

Conflict of interests: The manuscript was prepared and reviewed with the participation of the authors, who declare that there exists no conflict of interest that puts at risk the validity of the presented results.

BIBLIOGRAPHIC REFERENCES

Aloni, B., R. Cohen, L. Karni, L.H. Aktas, and M. Edelstein. 2010. Hormonal signaling in rootstock–scion interactions. *Sci. Hortic.* 127, 119-126. Doi: <https://doi.org/10.1016/j.scienta.2010.09.003>

Bhatt, R.M., K.K. Upreti, M.H. Divya, S. Bhat, C.B. Pavithra, and A.T. Sadashiva. 2015. Interspecific grafting to enhance physiological resilience to flooding stress in tomato (*Solanum lycopersicum* L.). *Sci. Hortic.* 182, 8-17. Doi: <https://doi.org/10.1016/j.scienta.2014.10.043>

Djidonou, D., A.H. Simonne, K.E. Koch, J.K. Brecht, and X. Zhao. 2016. Nutritional quality of field-grown tomato fruit as affected by grafting with interspecific hybrid rootstocks. *HortScience* 51, 1618-1624. Doi: <https://doi.org/10.21273/HORTSCI11275-16>

Feller, C., H. Bleiholder, M. Hess, U. Meier, T. Van Den Boom, D.L. Peter, L. Buhr, H. Hack, R. Klose, R. Stauss, E. Weber, and M. Philipp. 1997. Compendium of growth stage identification keys for mono and dicotyledonous plants extended BBCH scale. 2nd ed. In: <https://www.hortiadvice.dk/upl/website/bbch-skala/scaleBBCH.pdf>; consulted: March, 2020.

Fullana-Pericàs, M., J. Ponce, M.À. Conesa, A. Juan, M. Ribas-Carbó, and J. Galmés. 2018. Changes in yield, growth and photosynthesis in a drought-adapted Mediterranean tomato landrace (*Solanum lycopersicum* ‘Ramellet’) when grafted onto commercial rootstocks and *Solanum pimpinellifolium*. *Sci. Hortic.* 233, 70-77. Doi: <https://doi.org/10.1016/j.scienta.2018.01.045>

Gaion, L.A., L.T. Braz, and R.F. Carvalho. 2018. Grafting in vegetable crops: A great technique for agriculture. *Int. J. Veg. Sci.* 24, 85-102. Doi: <https://doi.org/10.1080/19315260.2017.1357062>

Geboloğlu, N., E. Yilmaz, P. Cakm. M. Aydin, and Y. Kasap. 2011. Determining of the yield, quality and nutrient content of tomatoes grafted on different rootstocks in soilless culture. *Sci. Res. Essays* 6, 2147-2153. Doi: <https://doi.org/10.5897/SRE10.1079>

Hossain, M.G., M.A. Ali, R.A. Ripa, S. Aydin, and S. Mahmood. 2019. Influence of rootstocks on yield and quality of summer tomato cv. ‘BARI Tomato-4’. *Earth Syst. Environ.* 3, 289-300. Doi: <https://doi.org/10.1007/s41748-019-00101-4>

Khah, E.M., E. Kakava, A. Mavromatis, D. Chachalis, and C. Goulas. 2006. Effect of grafting on growth and yield of tomato (*Lycopersicon esculentum* Mill.) in greenhouse and open-field. *J. Appl. Hortic.* 8, 3-7. Doi: <https://doi.org/10.37855/jah.2006.v08i01.01>

Kumar, P., Y. Rouphael, M. Cardarelli, and G. Colla. 2017. Vegetable grafting as a tool to improve drought resistance and water use efficiency. *Front. Plant Sci.* 8, 1130. Doi: <https://doi.org/10.3389/fpls.2017.01130>

Kumar, A.B., and K. Sanket. 2017. Grafting of vegetable crops as a tool to improve yield and tolerance against diseases. A review. *Int. J. Agric. Sci.* 9, 4050-4056.

Lee, J.M., C. Kubota, S.J. Tsao, Z. Bie, P.H. Echevarria, L. Morra, and M. Oda. 2010. Current status of vegetable grafting: Diffusion, grafting techniques, automation. *Sci. Hortic.* 127, 93-105. Doi: <https://doi.org/10.1016/j.scienta.2010.08.003>

Martínez-Andújar, C., J.M. Ruiz-Lozano, I.C. Dodd, A. Albacete, and F. Pérez-Alfocea. 2017. Hormonal and nutritional features in contrasting rootstock-mediated tomato growth under low-phosphorus nutrition. *Front. Plant Sci.* 8, 13. Doi: <https://doi.org/10.3389/fpls.2017.00533>

Milenković, L., J. Mastilović, Ž. Kevrešan, A. Bajić, A. Gledić, L. Stanojević, D. Cvetković, L.J. Šunić, and Z.S. Ilić. 2020. Effect of shading and grafting on yield and quality of tomato. *J. Sci. Food Agric.* 100, 623-633. Doi: <https://doi.org/10.1002/jsfa.10057>

Miskovic, A., O. Ilic, J. Bacanovic, V. Vujasinovic, and B. Kukic. 2016. Effect of eggplant rootstock on yield and quality parameters of grafted tomato. *Acta Sci. Polon. Hortic. Cul.* 15, 149-159.

Muneer, S., H. Ch. Ko, H. Wei, Y. Chen, and B.R. Jeong. 2016. Physiological and proteomic investigations to study the response of tomato graft unions under temperature stress. *PLoS ONE*, 11, 23. Doi: <https://doi.org/10.1371/journal.pone.0157439>

Ntatsi, G., D. Savvas, H.P. Kläring, and D. Schwarz. 2014. Growth, yield, and metabolic responses of temperature-stressed tomato to grafting onto rootstocks differing in cold tolerance. *J. Am. Soc. Hortic. Sci.* 139, 230-243. Doi: <https://doi.org/10.21273/JASHS.139.2.230>

Pogonyi, A., Z. Pék, L. Helyes, and A. Lugasi. 2005. Effect of grafting on the tomato's yield, quality and main fruit components in spring forcing. *Acta Aliment.* 34, 453-462. Doi: <https://doi.org/10.1556/AAlim.34.2005.4.12>

Qaryouti, M.M., W. Qawasmi, H. Hamdan, and M. Edwan. 2007. Tomato fruit yield and quality as affected by grafting and growing system. *Acta Hortic.* 741, 199-206. Doi: <https://doi.org/10.17660/ActaHortic.2007.741.22>

R Core Team. 2017. R: a language and environment for statistical computing. In: Vienna: R Foundation for Statistical Computing, <https://cran.r-project.org>; consulted: March, 2020.

Rahmatian, A., M. Delshad, and R. Salehi. 2014. Effect of grafting on growth, yield and fruit quality of single and double stemmed tomato plants grown hydroponically. *Hortic. Environ. Biotechnol.* 55, 115-119. Doi: <https://doi.org/10.1007/s13580-014-0167-6>

Reddy, P.P. 2016. Grafted vegetables for management of soilborne pathogens. pp.83-97. In: Reddy, P.P. (Ed.). *Sustainable crop protection under protected cultivation*. Springer, Singapore. Doi: https://doi.org/10.1007/978-981-287-952-3_7

Riga, P. 2015. Effect of rootstock on growth, fruit production and quality of tomato plants grown under low temperature and light conditions. *Hortic. Environ. Biotechnol.* 56, 626-638. Doi: <https://doi.org/10.1007/s13580-015-0042-0>

Savvas, D., G.B. Öztekin, M. Tepecik, A.M. Ropokis, Y. Tüzel, G. Ntatsi, and D. Schwarz. 2017. Impact of grafting and rootstock on nutrient-to-water uptake ratios during the first month after planting of hydroponically grown tomato. *J. Hortic. Sci. Biotechnol.* 92, 294-302. Doi: <https://doi.org/10.1080/14620316.2016.1265903>

Sen, A., R. Chatterjee, P. Bhaisare, and S. Subba. 2018. Grafting as an alternate tool for biotic and abiotic tolerance with improved growth and production of solanaceous vegetables: Challenges and scopes in India. *Int. J. Curr. Microbiol. App. Sci.* 7, 121-135. Doi: <https://doi.org/10.20546/ijcmas.2018.701.014>

Singh, H., P. Kumar, S. Chaudhari, and M. Edelstein. 2017. Tomato grafting: A global perspective. *HortScience* 52, 1328-1336. Doi: <https://doi.org/10.21273/HORTSCI11996-17>

Singh, H., P. Kumar, A. Kumar, M.C. Kyriacou, G. Colla, and Y. Rouphael. 2020. Grafting tomato as a tool to improve salt tolerance. *Agronomy* 10, 21. Doi: <https://doi.org/10.3390/agronomy10020263>

Torres, P.A. 2017. Tomate al aire libre. *Boletín INIA* 376. Instituto de Investigaciones Agropecuarias, Santiago.

Turhan, A., N. Ozmen, M.S. Serbeci, and V. Seniz. 2011. Effects of grafting on different rootstocks on tomato fruit yield and quality. *Hortic. Sci.* 38, 142-149. Doi: <https://doi.org/10.17221/51/2011-HORTSCI>