

Report and diagnoses of *Hambletonia pseudococcina* Compere (Hymenoptera: Encyrtidae) and *Pseudiasata* sp. (Diptera: Drosophilidae): Natural enemies of *Dysmicoccus brevipes* (Cockerell) (Hemiptera: Pseudococcidae) in Valle del Cauca, Colombia

Reporte y diagnosis de *Hambletonia pseudococcina* Compere (Hymenoptera: Encyrtidae) y *Pseudiasata* sp. (Diptera: Drosophilidae): Enemigos naturales de *Dysmicoccus brevipes* (Cockerell) (Hemiptera: Pseudococcidae) en el Valle del Cauca, Colombia



TAKUMASA KONDO^{1, 3}
JAMES B. WOOLLEY²
YENIFER CAMPOS-PATIÑO¹


An aggregation of pink pineapple mealybug, *Dysmicoccus brevipes* (Cockerell), with some parasitized mealybugs (green arrows) and mummies (yellow arrows) by *Hambletonia pseudococcina* Compere.

Photo: T. Kondo.

ABSTRACT

During a study carried out to determine the prevalence of mealybugs in pineapple (*Ananas comosus*) orchards conducted in the department of Valle del Cauca, Colombia, the parasitoid wasp *Hambletonia pseudococcina* Compere, 1936 (Hymenoptera: Encyrtidae) and the predatory fly *Pseudiasata* sp. (Diptera: Drosophilidae) were found in association with the pineapple mealybug *Dysmicoccus brevipes* (Cockerell, 1893) (Hemiptera: Pseudococcidae). These two natural enemies are briefly diagnosed and illustrated. This is the first record of the genus *Pseudiasata* in Colombia.

Additional key words: *Ananas comosus*; biological control; endoparasitoid; odoriferous secretions; species diagnosis; taxonomy.



¹ Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de Investigación Palmira, Palmira (Colombia). ORCID Kondo, T.: 0000-0003-3192-329X; ORCID Campos-Patiño, Y.: 0000-0002-5209-8340

² Texas A&M University, Department of Entomology, College Station (USA). ORCID Woolley, J.B.: 0000-0001-9893-2719

³ Corresponding author. tkondo@agrosavia.co

RESUMEN

Durante un estudio realizado para determinar la prevalencia de cochinillas harinosas en huertos de piña (*Ananas comosus*) realizado en el departamento del Valle del Cauca, Colombia, la avispa parasitoide, *Hambletonia pseudococcina* Compere, 1936 (Hymenoptera: Encyrtidae) y la mosca depredadora *Pseudiasata* sp. (Diptera: Drosophilidae) se encontraron en asociación con la cochinilla harinosa de la piña, *Dysmicoccus brevipes* (Cockerell, 1893) (Hemiptera: Pseudococcidae). Los dos enemigos naturales se diagnostican e ilustran brevemente. Este es el primer registro del género *Pseudiasata* en Colombia.

Palabras clave adicionales: *Ananas comosus*; control biológico; endoparasitoide; secreciones odoríferas; diagnosis de especies; taxonomía.

Received: 14-07-2021 Accepted: 11-09-2021 Published: 14-02-2022

INTRODUCTION

Pineapple, *Ananas comosus* Merr. (Bromeliaceae), is native to southern Brazil and Paraguay where it was domesticated and then was taken northward through South and Central America to Mexico and the West Indies by the indigenous people prior to the arrival of Europeans (Morton, 1987). Currently, pineapple is produced in tropical and subtropical regions around the world. According to FAOSTAT (2021), based on the average production (in metric tons) between 2014-2019, the top 10 pineapple producers worldwide are Brazil (2,193,653 t), Thailand (2,139,920 t), Philippines (2,042,208 t), Costa Rica (1,773,360 t), India (1,373,663 t), Indonesia (1,135,412 t), Nigeria (1,126,262 t), mainland China (994,521 t), Mexico (660,171 t), and Colombia (506,365 t). Based on the figures of the National Agricultural Survey, in 2015, there were a total of 8,871 ha planted for the cultivation of pineapple in Colombia, of which the department of Valle del Cauca reported the highest production with 35.22%, followed by Quindío with 25.29%, Santander 11.68%, Cauca 10.82% and Casanare with 7.20%; the remaining 9.79% were produced by the other 12 departments with shares each below 2.49%, such as Antioquia, Bolívar, Boyacá, Cesar, Córdoba, Cundinamarca, Huila, Meta, Nariño, Risaralda, Sucre, and Tolima (DANE, 2016).

Various insect pests have been reported affecting pineapple crops in Colombia, including the weevils *Metamasius dimidiatipennis* (Jekel) and *M. hebetatus* (Gyllenhal) (Coleoptera: Curculionidae: Dryophthorinae), the pink pineapple mealybug *Dysmicoccus brevipes* (Cockerell, 1893) (Hemiptera: Pseudococcidae), the pineapple hairstreak *Strymon*

megarus (Godart) (as *Strymon basilides*) (Lepidoptera: Lycaenidae), the sugarcane midge *Elaphria nucicolora* (Guenée, 1852) (Lepidoptera: Noctuidae), and the pineapple fly *Melanoloma viatrix* Hendel, 1911 (Diptera: Richardiidae) (Villalobos-Moreno *et al.*, 2009; Mosquera, 2016). Of these, in the department of Valle del Cauca, *M. dimidiatipennis*, *S. megarus* and *D. brevipes* are considered the main insect pests of pineapple (Mesa-Cobo *et al.*, 2014).

During a survey carried out in 2019 to determine the prevalence of mealybugs associated with two agroclimatic zones of pineapple production in the department of Valle del Cauca, Colombia, two natural enemies were collected which were identified as the encyrtid wasp *Hambletonia pseudococcina* Compere, 1936 (Hymenoptera: Encyrtidae) and the drosophilid fly *Pseudiasata* sp. (Diptera: Drosophilidae).

According to the Universal Chalcidoidea Database (Noyes, 2019), *H. pseudococcina* is widespread in the New World, recorded from the Antilles, Argentina, Bermuda, Brazil (Goias, Rondonia, Sao Paulo), Colombia, Costa Rica, Dominica, Dominican Republic, Ecuador, Jamaica, Mexico, Puerto Rico, Trinidad and Tobago, United States of America (California, Florida), and Venezuela. Outside the Americas, it is known from Africa (Ghana), the South Pacific region (Hawaii) and the Palaearctic region (Japan, Taiwan) (Noyes, 2019). Although not recorded in the Chalcidoidea database, *H. pseudococcina* has been recorded also in Cuba (Ceballos and Granada, 2004). *Hambletonia pseudococcina* has been recorded previously in Colombia from the departments of Caqueta reared

from *Pseudococcus* sp. (likely *Dysmicoccus*) (Hemiptera: Pseudococcidae) on pineapple (Sharkov and Woolley, 1997), and Valle del Cauca, from *Puto barberi* (Cockerell, 1895) (Hemiptera: Putoidae) on coffee roots (Gil *et al.*, 2016). This is the first report of *H. pseudococcina* from *D. brevipes* on pineapple in the department of Valle del Cauca.

The genus *Pseudiasata* Coquillett (Diptera: Drosophilidae) is New World in origin, known from the Mid-Atlantic region of the United States, down to Central and South America (Brake and Bächli, 2008). *Pseudiasata* includes six species, namely *P. armata* Wheeler, *P. australis* Blanchard, *P. brasiliensis* Costa Lima, *P. nebulosa* Coquillett, *P. pseudococcivora* Sabrosky, and *P. vorax* Sabrosky (Brake and Bächli, 2008). All known species of *Pseudiasata*, with the exception of *P. nebulosa*, the type species of the genus, and *P. armata*, have been recorded preying exclusively on pineapple mealybugs (Hardy, 1959). The insect hosts of *P. armata* and *P. nebulosa* are unknown. This is the first record of the genus in Colombia.

Here we document the first report of the parasitoid wasp *H. pseudococcina* and the predatory fly *Pseudiasata* sp., associated with the mealybug *D. brevipes*, in the department of Valle del Cauca and provide brief diagnoses for the recognition of these natural enemies.

MATERIALS AND METHODS

Collecting locality

Both the parasitoid wasp and the predatory fly were collected in association with pineapple mealybugs, which were found at the base of the crown of a pineapple fruit collected at Las Veraneras farm, municipality of Vijes, department of Valle del Cauca, Colombia (03°05'26" N, 76°28'53" W, 1,455 m a.s.l.). The infested fruit was found at the edge of a newly planted batch of pineapple crop within a pile of discarded plants from the previous crop.

Sampling and specimen preservation

The pineapple skin with the mealybug infestation and associated natural enemies was cut out from the fruit with a pocketknife, placed into a 200 mL plastic container and taken to the laboratory. Then the samples were transferred to a different plastic container

with a dry paper towel on the bottom and sealed with a mesh lid to allow air flow. The drosophilid fly pupae were found near the mealybugs and were kept in the same plastic container with the parasitized mealybugs until they emerged. The parasitized mealybugs and fly pupae were observed daily, and the emerged parasitoids and flies were placed in vials with 75% ethanol (EtOH). A few mealybug mummies (= dead parasitized mealybugs that become brownish to reddish brown in color) were dissected in order to photograph the developing parasitoids. Some samples of the encyrtid parasitoids were sent to the second author (J.B.W.) for identification.

Mass rearing of *Hambletonia pseudococcina*

Several specimens of *D. brevipes* parasitized by *H. pseudococcina* were taken to the entomology laboratory at the Corporacion Colombiana de Investigación Agropecuaria – Agrosavia, Palmira Research Station, located in the municipality of Palmira, Valle del Cauca, Colombia, 03°31'17" N, 76°18'25" W, ca. 1,000 m a.s.l., to start a rearing colony on mealybugs of *D. brevipes* kept on pumpkin, *Cucurbita moschata* Duchesne (Cucurbitaceae). The insects were reared under laboratory conditions (25.9 ± 1.7°C, with 67.8 ± 8.1% RH, and natural light regimen, approximately 12:12 h [L:D]). The pumpkins were stored in a plastic container with a mesh lid, with a dry paper towel on the bottom. The reared parasitoids were used for observations and for providing voucher specimens.

Specimen preparation and identification

Encyrtid wasp

The emerged parasitoids from the sampled mealybugs were preserved in 75% ethanol (EtOH). Some alcohol-preserved specimens were critical-point-dried and mounted following the method described by Noyes (1982). Some specimens were point-mounted without previous treatment and others were mounted on slides in Hoyer's mounting medium (distilled water 50 cm³, gum Arabic 30 cm³, chloral hydrate 200 cm³ and glycerin 20 cm³). The antennae were removed prior to the treatment of the specimens and used to determine the distribution pattern of the coloration. The specimens used for slide-mounting were cleared by heating them ca. 15 min in 10% w/v KOH prior to mounting. The specimens were identified to genus and species level by using the keys of Noyes

(2000) and Sharkov and Woolley (1997). Morphological terminology used for the parasitoids follows that of Gibson (1997).

Drosophilid fly

The external morphology of the drosophilid flies kept in 75% ethanol was observed under a Nikon SMZ 745T stereomicroscope. The specimens were identified to the genus level using the diagnoses provide by Hardy (1959) and Sabrosky (1951). Only two adult females emerged, and because males are critical for the taxonomy of these flies, species level identification was not possible. Sabrosky (1951) studied the adults of four species, namely *Pseudiasata nebulosa*, *P. pseudococcivora*, *P. vorax*, and an undescribed species, which he listed as *Pseudiasata* sp., and concluded that they were all virtually indistinguishable based on superficial characters. He pointed out that the differences in the wing markings were not useful for the separation of species because they were all very similar, differing only in extent and intensity, and that the fundamental pattern of the above species was the same (Sabrosky, 1951). Sabrosky (1951) also stated out that species separation of the adult stage could be done by comparing the lateral and anterior aspects of the male terminalia and that females can be identified only by association with male specimens.

Mealybugs

The mealybug specimens were slide-mounted following the methods of Williams and Granara de Willink (1992). The specimens were identified to species level using the keys of Caballero *et al.* (2017), Pacheco da Silva *et al.* (2020), Tanaka and Kamitani (2021) and Williams and Granara de Willink (1992).

Collecting permit

Insect samples were collected under a global collecting permit "Permiso marco de recolección de especímenes de especies silvestres de la diversidad biológica con fines de investigación científica no comercial" [Permit framework for collecting of specimens of wild species of the biological diversity for non-commercial scientific research purposes], resolution No. 1466, expedited on December 3, 2014, by the Autoridad Nacional de Licencias Ambientales (ANLA) [Colombian National Authority Environmental Permits].

Depositories

CTNI: Colección Taxonómica Nacional de Insectos "Luis María Murillo", Corporación Colombiana de Investigación Agropecuaria, Agrosavia, Mosquera (Colombia).

TAMUIC: Texas A&M University Insect Collection, Department of Entomology, Texas A&M University, College Station (TX, USA).

RESULTS AND DISCUSSION

Diagnoses

Hambletonia Compere is a distinctive genus in the subfamily Tetracneminae of Encyrtidae. It can be recognized by an orange or brown coloration of head and mesosoma (Fig. 1A), head strongly triangular in lateral aspect with a sharp occipital margin (Fig. 1B), and a sharply margined frontofacial ridge that continues ventrally as sharply margined antennal scrobes enclosing a concave facial area (Fig. 1C) (Noyes, 2000). Noyes (2000) diagnoses the subfamily Tetracneminae providing characteristics to distinguish them from the subfamily Encyrtinae. Sharkov and Woolley (1997) gave a detailed description of the genus *Hambletonia* and provided a key to world species.

Hambletonia pseudococcina Compere, 1936

Hambletonia pseudococcina can be diagnosed by the following combination of characters: i) anterior edge of frontovertex in dorsal view (occipital plane oriented vertically) distinctly convex (Fig. 1B); ii) frontovertex without prominence and punctures, or at most with scattered small inconspicuous punctures (Figs. 1A and B); iii) frontovertex more or less smooth between anterior ocellus and frontofacial ridge with only minute piliferous punctures; iv) frontovertex and eyes completely hairless, or with extremely minute hairs visible only at higher magnification when viewed and illuminated at certain angle, with at most a few very short hairs present behind posterior ocelli; v) setae on eyes at most about half as long as diameter of facet, vi) anterior and posterior ocelli situated in the same plane (Fig. 1A); vii) scape mostly dark brown; viii) pedicel with a tuft of elongate hairs on dorsal side (Fig. 1A), ix) tuft of setae on pedicel composed of very slightly broadened, almost scale-like setae




Figure 1. *Hambletonia pseudococcina* Compere. **A.** Female, dorsal habitus. **B.** Female, lateral habitus. **C.** Female, front of head and antennae. **D.** Female, fore wing. Photos by J.B. Woolley.

(Fig. 1C); x) funicle 6-segmented; xi) clava entirely dark brown and less than 1.5X as long as broad with sensory area as long as dorsal surface of clava; xii) forewing length 2.0-2.5 times maximum width (Fig. 1D); xiii) basal cell of fore wing pilose, propodeal spiracle not reaching side of propodeum, its diameter less than diameter of mid basitarsus; longitudinal median basal groove on scutellum deep and about half as long as scutellum; and xiv) costal cell of forewing with several rows of setae dorsally (Fig. 1D) (Sharkov and Woolley, 1997).

Hambletonia pseudococcina is close to *H. pilosifrons* Sharkov and Woolley (1997), *H. roseni* Sharkov and Woolley (1997), and *H. calvifrons* Sharkov and Woolley (1997). It differs from *H. pilosifrons* and *H. roseni* by the completely naked frontovertex and eyes, with at most minute hairs that can be present behind the posterior ocelli [*H. pilosifrons* with frontovertex with translucent brownish semierect to erect hairs directed anteriorly and some posteriorly; and eyes with rather dense translucent hairs; and *H. roseni* with extremely minute hairs on the frontovertex and eyes].

It further differs from *H. roseni* by the pilose basal cell of the forewing [basal part of forewing almost hairless in *H. roseni*]. *H. pseudococcina* differs from *H. calvifrons* by the convex anterior edge of the frontovertex; the presence of several rows of setae on the dorsal surface of the costal cell of the forewing; scape dark colored and maximally broadened in the middle part; and eyes almost hairless [*H. calvifrons* can be diagnosed by the following character states: i) anterior edge of frontovertex almost straight; ii) costal cell of forewing with only a single row of setae; iii) scape slightly more elongated and light colored and maximally broadened in distal half; iv) scape entirely pale orange; v) clava with base pale orange and at least 1.8X as long as broad with sensory area less than 0.9X as long as dorsal margin of clava; vi) tuft of setae on pedicel composed of gradually tapering, bristle-like setae; vii) frontovertex and eyes with minute hairs; and viii) setae on eye nearly as long as diameter of facet].

In life, the dorsum of unparasitized pineapple mealybugs is pinkish, usually with well-developed lateral

waxy filaments (Fig. 2A); ventral derm is yellowish pink, with yellowish limbs (Fig. 2B). Mealybug mummies parasitized by *H. pseudococcina* are almost spherical (Figure 2C–H), with brown to reddish-brown limbs and derm, and have noticeably less mealy wax on their surfaces (Figure 2D–I) compared to unparasitized individuals and often have shorter lateral wax filaments (Fig. 2C). One hundred mummies were studied. Most mummies (83%) had one parasitoid, however some mummies (17%) showed super-parasitism, with two parasitoids within when dissected; those mummies that contained two individuals within (Fig. 2F and I) were larger and much broader than those with only one individual within (Fig. 2E). Fig. 2I shows a mummy that had two parasitoid pupae within, of which one was extracted (Fig. 2J). The parasitoid pupa was of the exarate type, whitish, semitranslucent, with red eyes.

A small but odoriferous parasitoid

While processing the samples of *H. pseudococcina*, Y.C.P. noticed that the adult wasps produce a strong distinctive smell, which persisted in the Petri dishes

even several weeks after they were removed. We asked eight persons (including Y.C.P. and T.K.) to describe the smell of *H. pseudococcina*, and their opinions were as follows: i) a sweet smell similar to coconut essence; ii) fermented wood; iii) cedar wood with a touch of honey; iv) a smell similar to that produced by dolichoderine ants; and v) wet cigarette butts, with a sweet honey scent. From all the above descriptions of the smell, the coconut essence smell was detected by five of them. This appears to be the first report of a distinctive smell associated with a member of the Encyrtidae. Further studies are needed to determine the composition of this odoriferous secretion and its purpose, e.g., whether it is a sex pheromone, a host marking pheromone, a deterrent for other parasitoids and predators, etc.

Pseudiasata Coquillett

The genus *Pseudiasata* can be diagnosed as follows. *Head*: eyes bare, large, occupying most of the area of head in profile; cheek linear; front with numerous short hairs; occiput distinctly concave; arista pubescent when seen under the microscope. *Thorax*: densely




Figure 2. Unparasitized and parasitized specimens of *D. brevipes* by *H. pseudococcina*. A. Dorsal view of unparasitized mealybug. B. Ventral view of unparasitized mealybug. C. Dorsal view of parasitized mealybug. D. Ventral view of parasitized mealybug. E. Sideview of parasitized mealybug with single parasitoid. F. Sideview of mealybug with super-parasitism. G. Dorsal view of parasitized mealybug in advance stage of parasitoid development. H. Ventral view of parasitized mealybug in advance stage of parasitoid development. I. Parasitized mealybug with two parasitoid pupae, one of which was removed. J. Pupa of *H. pseudococcina*. Photos: A and B by T. Kondo; C–I by Y. Campos-Patiño.



Figure 3. A. Aggregation of pineapple mealybugs near base of the fruit crown, with a pupa of *Pseudiastata* sp. on left side (see arrow). B. An empty pupa of *Pseudiastata* sp. C. Adult female *Pseudiastata* sp. Notice the characteristic color patterns on the wings. Photos: T. Kondo.

covered with hairs but not arranged in definite rows; scutellum bare but with bristles; mesopleuron and pteropleuron also bare; legs normal, short, preapical dorsal bristles present in at least some tibiae; wings each pictured (with markings), venation and color pattern distinct (Fig. 3), costa fractured in two places, with a strong bristle before each break, costa extending to fourth vein, discal and second basal cells merging, anal cell small. *Chaetotaxy*: 1 inner and 1 outer vertical, cruciate postvertical bristles short, cruciate ocellar bristles short, 3 pairs of fronto-orbital bristles (anterior pair convergent, the 2 posterior pairs reclinate), 1 humeral bristle, 1 presutural bristle, 1 + 1 notopleural bristle, 1 strong supraalar bristle, 2 postalar bristles, 2 posterior dorsocentral bristles, 1 prescutellar acrostichal bristle, 2 scutellar bristles, and 2 sternopleural bristles (Sabrosky, 1951).

The two *Pseudiastata* sp. specimens collected in the present study are each about 3 mm long. They match fairly well the generic diagnoses by Sabrosky (1951); however, further field studies are needed to collect male specimens of *Pseudiastata* in order to carry out a species level identification. A key to species of *Pseudiastata* can be found in Hardy (1959).

Material studied

Dysmicoccus brevipes (Cockerell) (Pseudococcidae), det. Takumasa Kondo, ex root crown of *Ananas comosus* L. cv. MD2 (Bromeliaceae), **Colombia**: Valle del Cauca, municipio Vijes, 03°45' N, 76°28' W, 1,455 m

a.s.l., 11.iv.2019, coll. Takumasa Kondo, 2 slides: 10 adult females (CTNI: Catalogue No. 6693).

Hambletonia pseudococcina Compere, det. J.B. Woolley, **Colombia**: Valle del Cauca, Municipio de Vijes, 03°45' N, 76°28' W, 1,455 m a.s.l., 11.iv.2019, ex *Dysmicoccus brevipes* Cockerell found on fruit of pineapple, *Ananas comosus* L. cv. MD-2, coll. Yenifer Campos-Patiño, 12 female specimens, (TAMUIC: voucher specimen #758); 60 specimens in 75% alcohol (CTNI: Catalogue No. 6881).

Pseudiastata sp. (Diptera: Drosophilidae), det. Takumasa Kondo, **Colombia**: Valle del Cauca, Municipio de Vijes, 1,455 m a.s.l., 11.iv.2019, 03°45' N, 76°28' W, found next to an aggregation of *Dysmicoccus brevipes* Cockerell, on fruit of pineapple, *Ananas comosus* L. cv. MD-2, coll. Y. Campos-Patiño, 2 adult female specimens (CTNI: Catalogue No. 6682).

CONCLUSIONS

Two natural enemies were found in the department of Valle del Cauca, Colombia, in association with the pineapple mealybug, *Dysmicoccus brevipes* (Hemiptera: Pseudococcidae): an encyrtid parasitoid wasp, *Hambletonia pseudococcina* (Hymenoptera: Encyrtidae) and a drosophilid predatory fly, *Pseudiastata* sp. (Diptera: Drosophilidae). The genus *Pseudiastata* is for the first time recorded from Colombia and an odoriferous secretion is for the first time reported in a member of the family Encyrtidae. Further studies

are needed to explore the rich biodiversity of natural enemies in Colombia which have the potential to be incorporated into biological control programs.

Conflict of interests: The manuscript was prepared and reviewed with the participation of the authors, who declare that there exists no conflict of interest that puts at risk the validity of the presented results.

ACKNOWLEDGMENTS

The authors thank Dr. Penny J. Gullan (Australian National University, Canberra) and two anonymous reviewers for kindly reviewing the manuscript. This research Project was funded in part by the Colombian Corporation for Agricultural Research – Agrosavia, project: “Estrategias para el manejo integrado de las principales enfermedades y plagas que se presentan en el cultivo de piña (cultivar MD2) en los departamentos del Valle del Cauca y Santander”, Grant No. TV19.

BIBLIOGRAPHIC REFERENCES

- Brake, I. and G. Bächli. 2008. World catalogue of insects in Drosophilidae (Diptera), Vol. 9. Apollo Books, Stensrup, Denmark.
- Caballero, A., A.A. Ramos-Portilla, and T. Kondo. 2017. Scale insects (Hemiptera: Coccoomorpha) on sugarcane in Colombia, with description of a new species of *Tillanococcus* Ben-Dov (Coccidae). *Zootaxa* 4258(5), 490-500. Doi: 10.11646/zootaxa.4258.5.6
- Ceballos, M.R. and C. Granada. 2004. *Hambletonia pseudo-coccina* Compere (Hymenoptera: Chalcidoidea: Encyrtidae): New report in Cuba. *Rev. Prot. Veg.* 19(1), 70.
- DANE, Dirección Nacional de Estadística Colombia. 2016. Principales características del cultivo de la piña (*Ananas comosus* L.). Insumos y factores asociados a la producción agropecuaria. Boletín Mensual No. 54. https://www.dane.gov.co/files/investigaciones/agropecuario/sipsa/Bol_Insumos_dic_2016.pdf; consulted: June 2021.
- FAOSTAT. 2021. FAOSTAT database. In: <http://www.fao.org/faostat/en/#data/QC/visualize>; consulted: June, 2021.
- Gibson, G.A.P. 1997. Morphology and terminology. pp. 16-44. In: Gibson, G.A.P., J.T. Huber, and J.B. Woolley (eds.). Annotated keys to the genera of Nearctic Chalcidoidea. NRC Research Press, Ottawa, Canada.
- Gil, P.Z.N., M.P. Benavides, and L.M. Constantino-C. 2016. Hemiptera: Coccoidea de las raíces del café y sus parasitoides en el Valle del Cauca. *Rev. Cenicafe* 67(1), 73-80.
- Hardy, D.E. 1959. A review of the genus *Pseudiastata* Coquillett (Drosophilidae, Diptera). *Proc. Hawaii Entomol. Soc.* 17(1), 76-82.
- Mesa-Cobo, N., E.D. Gómez, C. Huertas, Y.M. Mena, C.A. Hernandez, E.D. Henao, and M. Cobo. 2014. Programa de manejo fitosanitario en piña en el departamento del Valle del Cauca. Universidad Nacional de Colombia, Sede Palmira, Facultad de Ciencias Agropecuarias, Palmira, Colombia.
- Morton, J. 1987. Pineapple. pp. 18-28. In: Fruits of warm climates. Julia F. Moton, Miami, FL. In: <https://hort.purdue.edu/newcrop/morton/pineapple.html>; consulted: June 2021.
- Mosquera Ramírez, S.Y. 2016. Identificación y caracterización de plagas y enfermedades asociadas con el cultivo tradicional de piña *Ananas comosus* (L. Merr) en comunidades del municipio de Lloró, departamento del Chocó, Colombia. *Rev. Bioetnia* 13(1), 78-86. Doi: 10.51641/bioetnia.v13i1.163
- Noyes, J.S. 1982. Collecting and preserving chalcid wasp (Hymenoptera: Chalcidoidea). *J. Nat. Hist.* 16(3), 315-334. Doi: 10.1080/00222938200770261
- Noyes, J.S. 2000. Encyrtidae of Costa Rica (Hymenoptera: Chalcidoidea), 1, The Subfamily Tetracneminae, Parasitoids of Mealybugs (Homoptera: Pseudococcidae). pp. 1-355. In: *Memoirs of the American Entomological Institute*. Vol. 62. Gainesville, FL.
- Noyes, J.S. 2019. Universal Chalcidoidea database. In: <http://www.nhm.ac.uk/chalcidoids/>; consulted: March, 2019.
- Pacheco da Silva, V.C., M.B. Kaydan, and C. Basso. 2020. Pseudococcidae (Hemiptera: Coccoomorpha) in Uruguay: morphological identification and molecular characterization, with descriptions of two new species. *Zootaxa* 4894(4), 501-520. Doi: 10.11646/zootaxa.4894.4.1
- Sabrosky, C.W. 1951. Two new species of *Pseudiastata* (Dipt., Drosophilidae) predacious on the pineapple mealybug. *Bull. Entomol. Res.* 41(3), 623-627. Doi: 10.1017/S0007485300027875
- Sharkov, A. and J.B. Woolley. 1997. A revision of the genus *Hambletonia* Compere (Hymenoptera: Encyrtidae). *J. Hymenopt. Res.* 6(2), 191-218.
- Tanaka, H. and S. Kamitani. 2021. Two new species of Coccoomorpha (Hemiptera: Sternorrhyncha) collected from Japanese silver grass, *Miscanthus sinensis* (Poaceae) in Okinawa Island, Japan. *Zootaxa* 4941(4), 569-579. Doi: 10.11646/zootaxa.4941.4.6
- Villalobos-Moreno, A., J.C. Agudelo-M., A.F. Barajas-S., S. Herrera-Q., and J.C. Céspedes. 2009. Cambios morfológicos durante la ontogenia del estado pupal de *Melanoloma viatrix* (Diptera: Richardiidae). *Rev. Colomb. Entomol.* 35(1), 98.
- Williams, D.J. and M.C. Granara de Willink. 1992. Mealybugs of Central and South America. CAB International, London.