

Assessment of the health status of citrus propagation material against *Hop stunt viroid* through biological and molecular diagnosis

Evaluación de la sanidad de material de propagación de cítricos frente al viroide del enanismo del lúpulo mediante diagnóstico biológico y molecular

DIANA MILENA RODRÍGUEZ-MORA^{1,2}

KEVIN ALEJANDRO RODRÍGUEZ-ARÉVALO¹

MAURICIO FERNANDO MARTÍNEZ¹

NUBIA MURCIA-RIAÑO^{1,2}

¹ Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de investigación Palmira, Palmira, (Colombia). ORCID Rodríguez-Mora, D.M.: <https://orcid.org/0000-0001-6545-3746>; ORCID Rodríguez-Arévalo, K.A.: <https://orcid.org/0000-0002-1058-8464>; ORCID Martínez, M.F.: <https://orcid.org/0000-0002-8145-1764>; ORCID Murcia-Riaño N.: <https://orcid.org/0000-0002-5679-3885>

² Corresponding author. dmrodriguez@agrosavia.co; nmurcia@agrosavia.co



Collection of micrografted citrus varieties, preserved in an aphid-proof mesh greenhouse at Agrosavia, Centro de Investigación Palmira, Valle del Cauca, Colombia.

Photo: N. Murcia-Riaño

Last name: RODRÍGUEZ-MORA / RODRÍGUEZ-ARÉVALO / MARTINEZ / MURCIA-RIAÑO

Short title: ASSESSMENT OF CITRUS AGAINST *HOP STUNT VIROID*Doi: <https://doi.org/10.17584/rcch.2024v18i2.17106>

Received: 18-01-2024 Accepted: 02-05-2024 Published: 10-06-2024

ABSTRACT

Hop stunt viroid (HSVd) is the causal agent of citrus cachexia, which can lead to significant crop losses. As part of a preventive management strategy, it is essential to use healthy plants from a certification program. This study aimed to, healthiness of 17 citrus cultivars was evaluated against HSVd using both biological and molecular diagnostics. For biological diagnosis, Etrog citron (*Citrus medica* L.) clone Arizona 861 S1 was used as an indicator plant, incorporating negative and positive controls for HSVd infection. For molecular diagnosis, reverse transcription polymerase chain reaction (RT-PCR) technique was employed with specific primers for viroid detection. Results indicated that six months after inoculation, the positive control exhibited symptoms associated with viroid infection, such as necrosis in the petiole and stem, leaf epinasty, and necrosis in the midvein. In contrast, both the negative control and evaluated citrus cultivars showed no symptoms. These findings were supported by molecular analysis, which allowed the amplification of a viroid genome region only in the positive control. Both biological and molecular diagnosis confirmed the absence of HSVd in the citrus varieties assessed. This study will facilitate the evaluation of the citrus plant collections used in the foundation category as a source of buds for plant multiplication of registered and certified seeds for citrus propagation nurseries in Colombia.

Additional key words: viroid; bioamplifier; RT-PCR; citrus; diagnosis; indexing.

RESUMEN

El viroide del enanismo del lúpulo (HSVd) es el agente causal de la cachexia de los cítricos, que puede provocar pérdidas significativas en el cultivo. Como parte de una estrategias de manejo preventivo, es indispensable utilizar plantas sanas provenientes de un programa de certificación. Este estudio tuvo como objetivo evaluar la sanidad de 17 cultivares de cítricos frente a HSVd, utilizando diagnósticos biológicos y moleculares. Para el diagnóstico biológico se empleó como

planta indicadora cidro Etrog (*Citrus medica* L.) clon Arizona 861 S1, incluyendo controles negativos y positivos para la infección por HSVd. Para el diagnóstico molecular se utilizó la técnica de reacción en cadena de la polimerasa con transcriptasa inversa (RT-PCR) con cebadores específicos para la detección del viroide. Los resultados indicaron que seis meses después de la inoculación, el control positivo mostró síntomas asociados con la infección por el viroide, como necrosis en el peciolo y tallo, epinastia foliar y necrosis en vena media. Por el contrario, tanto el control negativo como los cultivares de cítricos evaluados no presentaron síntomas. Estos resultados fueron corroborados por el análisis molecular, que permitieron amplificar una región del genoma del viroide, únicamente en el control positivo. Ambos diagnósticos biológico y molecular confirmaron la ausencia de HSVd en las variedades de cítricos evaluadas. Este estudio facilitará evaluación de las colecciones de plantas de cítricos utilizadas en la categoría fundación como fuente de yemas para la multiplicación de semilla registrada y certificada para los viveros productores de material de propagación de cítricos en Colombia.

Palabras clave adicionales: viroide; amplificador biológico; RT-PCR; cítricos; diagnóstico; indexaje.

INTRODUCTION

Colombia has an approximate cultivated area of 87,638 ha of citrus, encompassing orange, mandarin, and acid lime crops (MADR, 2021). However, its production faces various technological limitations, with one of the most significant challenges being the scarce availability of high-quality planting material, free from graft-transmitted diseases.

Various pathogens causing systemic diseases threaten citrus production, including *Citrus tristeza virus* (CTV) (*Closterovirus: Closteroviridae*) (Dawson *et al.*, 2015), *Candidatus Liberibacter asiaticus* (*Candidatus Liberibacter: Rhizobiaceae*) (Bové, 2006) *Citrus exocortis viroid* (CEVd) (*Pospiviroid: Pospiviroidae*) (Semancik and Weathers, 1972), *Citrus dwarfing viroid* (CDVd) (*Apscaviroid: Pospiviroidae*) (Rakowski *et al.*, 1994) y *Hop stunt viroid* (HSVd) (*Hostuviroid: Pospiviroidae*) (Ohno *et al.*, 1983). All these pathogens have been reported in Colombia (Murcia *et al.*, 2010; Murcia *et al.*, 2020).

Among viroid-caused diseases, cachexia, formerly known as xyloporosis, stands out, with its causal agent being HSVd (Serra *et al.*, 2018). Viroids are infectious agents consisting of circular RNA, single-stranded molecules without coat protein, of variable length ranging from 250 to 400

nucleotides (Venkataraman *et al.*, 2021). Viroid replication depends on factors and enzymes encoded by the host since they lack the cellular machinery necessary for this process (Serra *et al.*, 2018). Biologically, HSVd has been classified into pathogenic variants type CVd-IIb and CVd-IIc (Serra *et al.*, 2008), capable of inducing cachexia symptoms, and non-pathogenic variants of type CVd-IIa, which infect citrus plants without inducing symptoms (Semancik *et al.*, 1988). It has been established that the type and severity of citrus cachexia symptoms depend on co-infections with other citrus viroids in the tree (Vernière *et al.*, 2004). In susceptible hosts, a negative impact on yield has been observed, with reductions ranging from 34 to 76% and a decrease in fruit diameter (Belabess *et al.*, 2021). The primary mode of spread is through grafting with buds infected with the viroid and the use of non-disinfected pruning tools. No vectors are known to be involved in the transmission of this disease (Duran-Vila, 2004; Barbosa *et al.*, 2005).

Cases of cachexia affecting various citrus species have been reported worldwide, including Iran (Teymuri *et al.*, 2022; Zeitouni *et al.*, 2023), China (Wang *et al.*, 2010), the United States (Reanwarakorn and Semancik, 1999), and Brazil (Eiras *et al.*, 2013). In Colombia, HSVd infection has been detected in Tahiti acid lime (*Citrus x latifolia* Tan.) in open field nurseries and commercial orchards (Mosquera *et al.*, 2015; Rodríguez-Mora *et al.*, 2015; Murcia *et al.*, 2010). However, all identified isolates lacked the cachexia expression motif, indicating that they were non-pathogenic variants (Murcia *et al.*, 2010; Mosquera *et al.*, 2015).

Citrus cultivars affected by cachexia may exhibit visible symptoms or be asymptomatic (Vamenani *et al.*, 2014; Belabess *et al.*, 2021). In susceptible hosts such as Tangelo (*C. reticulata* Bla. × *C. paradisi* Macf.), symptoms include wood cracking, corresponding to wounds on the cambial face of the cortex, accompanied by discoloration of the phloem and gummy exudations (Duran-Vila *et al.*, 2000). In advanced stages of the disease, plants may develop stem pitting on the trunk, as well as leaf chlorosis and stunting (Serra *et al.*, 2008). Under favorable conditions for symptom manifestation, this disease can conduct fatal consequences for plants (Reanwarakorn and Semancik, 1988). During the nursery stage, infected plants are asymptomatic, making it necessary to guarantee healthiness of planting material.

The detection of pathogens causing systemic diseases, including viroids, is carried out through biological methods and molecular techniques. For the biological diagnosis of citrus viroid diseases, the Arizona 861-S1 clonal Etrog citron (*C. medica* L.) is employed as an indicator plant,

as recommended by Camps *et al.* (2014). This method involves observing the indicator plant's responses to viroid infection. For molecular diagnosis, Reverse Transcription Polymerase Chain Reaction (RT-PCR) is used, as indicated by Guerrero *et al.* (2013). RT-PCR enables specific amplification of the viroid genome, providing a precise and efficient tool to confirm the presence of the infectious agent. The combination of biological and molecular methods offers a comprehensive approach to the diagnosis of systemic diseases, facilitating more precise and rapid viroid detection.

Managing viroid-induced diseases, such as citrus cachexia, requires the implementation of preventive measures, including the utilization of disease-free propagation material produced within a certification program that mandates compliance with molecular and biological diagnostics (NAPPO, 2013). Since 2019, Colombia has regulated the production of citrus plants in protected environments, classifying the multiplication of propagative material and seeds into four categories: 1) genetic, 2) foundation, 3) registered, and 4) certified (ICA, 2019). The foundation category represents propagative material obtained from genetic seeds, serving as the origin for the registered and certified seed categories. The Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de Investigación Palmira, maintains a collection of citrus varieties obtained through the micrografting method. These varieties are preserved under aphid-proof mesh greenhouse conditions and belong to the foundation seed category. Plants in this category are used as mother plants in the country's citrus certification program. The aim of this research was to assess the healthiness of citrus propagative material in the foundation category, comprising 17 citrus cultivars, against HSVd. This evaluation was conducted through both biological and molecular diagnosis, so that it can be used as a reliable source of buds intended for plant multiplication in the registered and certified seed category, contributing to maintaining quality and healthiness in citrus nurseries in Colombia.

MATERIALS AND METHODS

Plant material

The study was conducted in December 2021 at Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de Investigación Palmira, in the department of Valle del Cauca, Colombia. Within the center's facilities, 17 citrus varieties planted in pots have been

maintained under controlled conditions inside an aphid-proof mesh greenhouse. These varieties are part of the elite citrus collection, obtained through *in vitro* micrografting of shoot apices. The plant collection consists of tangelos, grapefruits or pomelos, mandarins, oranges, and acid limes (Tab. 1). Each variety is represented by four replicates, totaling 68 plants.

Table 1. Collection of 17 micrografted citrus varieties.

No	Common name	Scientific name
1	Minneola tangelo	<i>C. reticulata</i> Bla. x <i>C. paradisi</i> Macf.
2	Orlando tangelo	<i>C. reticulata</i> Bla. x <i>C. paradisi</i> Macf.
3	Clemenules mandarin	<i>C. clementina</i>
4	ICA Bolo mandarin	<i>C. reticulata</i> Bla.
5	Oneco mandarin	<i>C. reticulata</i> Bla.
6	Arrayana mandarin	<i>C. reticulata</i> Bla.
7	Campbell Valencia orange	<i>C. sinensis</i> (L.) Osb.
8	Frost Valencia orange	<i>C. sinensis</i> (L.) Osb.
9	García Valencia orange	<i>C. sinensis</i> (L.) Osb.
10	Olinda Valencia orange	<i>C. sinensis</i> (L.) Osb.
11	Salustiana orange	<i>C. sinensis</i> (L.) Osb.
12	Sweety orange	<i>C. sinensis</i> (L.) Osb.
13	Frost Washington orange	<i>C. sinensis</i> (L.) Osb.
14	Valle Washington orange	<i>C. sinensis</i> (L.) Osb.
15	Key lime	<i>C. x aurantiifolia</i> (Christm.) Swing.
16	Tahiti lime	<i>C. x latifolia</i> Tan.
17	Pomelo	<i>C. paradisi</i> Macf.

Biological diagnosis

For biological indexing, the Etrog citron (*C. medica* L.) clone Arizona 861-S1, grafted onto the rootstock Volkameriana (*C. volkameriana* Ten. and Pasq.), served as an indicator plant. When Etrog citron plants reached 25 cm (from the graft union point), they were inoculated through vascular tissue contact with cortex from the 68 plants comprising the elite citrus collection, corresponding to the 17 micrografted citrus varieties. A branch from each quadrant was taken for each citrus variety, and these branches were used to inoculate the Etrog citron plants with two cortex segments (Fig. 1). Four replicates of each citrus variety were inoculated into three Etrog citron plants. The experimental setup included negative controls, which consisted of three healthy non-inoculated Etrog citron plants obtained by micrografting of shoot apices, and positive control, which consisted of three healthy Etrog citron plants inoculated with an HSVd isolate obtained from the viroid strain collection of Agrosavia, conserved *in vivo* and previously confirmed by RT-PCR and sequencing (Mosquera *et al.*, 2015).




Figure 1. Biological indexing of 17 citrus varieties on Etrog Citron (*C. medica* L.). A) collection of micrografted citrus varieties, preserved in an aphid-proof mesh greenhouse at Agrosavia, Centro de Investigación Palmira. B) collection of young shoots from citrus

varieties. C) removal of the Etrog citron cortex for inoculation, and D) inoculation of Etrog citron with citrus cortex segments. E) Etrog citron plant inoculated with two cortex segments. Photo: Research Group - Agrosavia, Centro de Investigación Palmira.

The inoculated plants were placed in an automated mesh greenhouse to maintain suitable environmental conditions (temperature between 28 and 32°C) to promote viroid multiplication (Duran-Vila, 1989). Visual inspections were conducted every 15 d over 6 months to monitor the expression of disease symptoms.

Molecular diagnosis

Six months after inoculation (MAI), plant tissue was collected from Etrog citron plants inoculated with cortex segments from the 17 micrografted citrus varieties, tissue from the negative control (non-inoculated Etrog citron plant), and from the positive control infected with HSVd. Each sample consisted of a pool of three plants, and four leaves per plant were taken, totaling 70 samples (68 samples corresponding to the 17 citrus varieties, one negative control, and one positive control). The samples were packed in properly labeled paper bags and transported to the molecular genetic laboratory for processing.

RNA extraction

RNA extraction was performed using 100 mg of the leaf midrib tissue pulverized with liquid nitrogen, following the CTAB protocol described by Chang *et al.* (1993). To this tissue, 900 µL of CTAB extraction buffer was added [2% CTAB (Cetyl Trimethyl Ammonium Bromide), 2% PVP (polyvinylpyrrolidone), 100 mM Tris-HCl (pH 8), 25 mM EDTA, 2 M NaCl, 0.5 g L⁻¹ spermidine, 2% β-mercaptoethanol], and it was placed in a water bath at 65°C for 15 min. The samples were shaken every 5 min. 1 mL of chloroform: Isoamyl alcohol (24:1) was added, vortexed for 10 s, and centrifuged at 12,000 rpm for 15 min at 4°C. The supernatant was transferred to a new tube, and the process was repeated. The aqueous phase was transferred to a new tube, and 1 mL of 4 M LiCl was added. The samples were incubated overnight at 4°C. They were then centrifuged at 12,000 rpm for 30 min at 4 °C, the supernatant was discarded, and the pellet was resuspended in 500 µL of TE-SDS buffer at 37°C. 700 µL of isopropanol and 200 µL of 5M NaCl were added, the tube was inverted several times, and it was incubated at -20°C for 1 h. The sample was centrifuged at 12,000 rpm for 10 min at 4 °C, the supernatant was discarded,

and the pellet was resuspended in 500 μ L of 70% ethanol. It was centrifuged at 12,000 rpm for 15 min at 4°C, the supernatant was discarded, and the pellet was dried for 1 h at room temperature in a laminar flow cabinet. The pellet was resuspended in 30 μ L of RNase-free water and stored at -80°C. The quality and quantity of RNA in the samples were assessed using spectrophotometry with the NanoDrop™ 1000 (ThermoFisher), analyzing the 260/280 and 260/230 absorbance ratios.

RT-PCR

Complementary DNA (cDNA) synthesis was carried out following the instructions of the SuperScript® III First-Strand Synthesis System for RT-PCR kit, catalog No. 18080-051 (Invitrogen). A mixture of 210 ng of RNA, 5.5 ng of random primer, and 0.83 mM of dNTPs (total) was denatured at 95°C for 5 min, resulting in a final volume of 9 μ L. The reaction was placed on ice for 1 min, and 5 μ L of a mixture containing 2X RT Buffer, 10 mM MgCl₂, 20 mM DTT, 20 U RNaseOUT, and 100 U SuperScript™ III RT was added. The reaction was incubated at 60°C for 1 h. The reaction was terminated at 85°C for 5 min, placed on ice for 1 min, 1 U of *Escherichia coli* RNase H was added, and it was incubated at 37°C for 20 min. The resulting cDNA was stored at -20°C.

The quality of the cDNA was verified by amplifying the 18S ribosomal RNA (rRNA) region using the primers 18SF (5' GAGAAACGGCTACCACATCCA 3') and 18SR (5' CGTGCCATCCAAAGTCCAAC 3'), following the conditions described by Du *et al.* (2006) with some modifications: reaction setup consisted of 1X Taq buffer (KCl), 1.5 mM of MgCl₂, 0.2 mM of dNTPs mixture, 0.2 μ M of each primer, 0.5 U of Taq DNA polymerase (ThermoScientific) in a final volume of 12.5 μ L. For the detection of HSVd, the specific primer combination HSVd/F2 (5'- GTGGCATCACCTCTCGGTT -3') and HSVd/R1 (5'-GGGGCTCCTTTCTCAGGTAAGTC -3') was used, following the conditions specified by Bernad and Duran-Vila (2006) with some modification as follows: the reaction was setup with 1X Taq buffer (KCl), 1.5 mM of MgCl₂, 0.5 mM of dNTPs mixture, 0.2 μ M pf each primer, 0.5 U of Taq DNA polymerase (ThermoScientific) in a final volume of 12.5 μ L.

Amplification process was carried out in an Agilent technologies Sure Cyclor 8800 with the following programs: for 18S amplification an initial denaturation of 95°C for 5 min was followed by 35 cycles of 95°C for 30 s, 52°C for 60 s and 72°C for 60 s, with a final elongation step of

72°C for 10 min and a cool-down to 20°C for 5 min. For HSVd/F2-R1, an initial denaturation at 94°C for 3 min was followed by 35 cycles of 94°C for 30 s, 60°C for 30 s and 72°C for 45 s, with a final elongation step of 72°C for 10 min and a cool-down to 20°C for 5 min. The amplified products were visualized on a 1.6% (w/v) agarose gel, run in 0.5X TBE buffer at 100 V, 400 mA for 80 min. Nucleic acids were stained with GelRed® (Biotium) at a concentration of 1.3X added to the PCR products, and visualization was done using the Enduro GDS gel documentation system (Labnet).

RESULTS AND DISCUSSION

Biological diagnosis

Indicator plants of Etrog citron clone Arizona 861-S1, inoculated with cortex tissue segments from the 17 micrografted citrus varieties were compared with reference controls: the negative control (non-inoculated Etrog citron plant) and the positive control infected with the HSVd viroid. This method allowed the expression of symptoms in the host when the viroid infection occurred.

At 2 MAI, initial symptoms of HSVd infection were observed in the three inoculated replicates of positive control, characterized by petiole necrosis. At 4 MAI, petiole necrosis and leaf epinasty were evident. After 5 months, symptoms progressed to include petiole necrosis, stem necrosis, leaf epinasty, and basal and middle necrosis of the central leaf vein. Finally, at 6 MAI, a more advanced stage was reached, with leaf epinasty (Fig. 2A), petiole necrosis (Fig. 2B), stem necrosis (Fig. 2C), and necrosis of the central leaf vein at the basal, middle, and apical parts observed (Fig. 2D). These symptoms have been associated with infections caused by viroids in Etrog citron (Duran-Vila *et al.*, 1988; Camps *et al.*, 2014; Zeitooni *et al.*, 2023). The plants used as negative controls (non-infected plants) (Fig. 2E, F, G, H) and Etrog citron plants inoculated with tissue segments from the 17 citrus varieties corresponding to the foundation seed (Fig. 2I, J, K, L) did not show any symptoms during the evaluation period. The presence of symptoms in positive controls, along with their absence in negative controls, demonstrates the quality of the biological indexing setup.




Figure 2. Biological indexing of HSVd in Etrog citron (*C. medica*) clone Arizona 861 S1, 6 months after inoculation. Positive control: A, B, C and D. Negative control: E, F, G and H. Inoculation of tissue from Minneola tangelo: I, J, K and L. Photo: Research group-Agrosavia, Centro de Investigación Palmira.

Etrog citron clone Arizona 861-S1 is considered one of the best hosts for conducting biological diagnosis of viroids due to its high susceptibility and symptom expression for most viroids reported in citrus (Duran-Vila *et al.*, 1988; Villalobos *et al.*, 2004; Camps *et al.*, 2014; Murcia *et al.*, 2020; Rodríguez-Mora *et al.*, 2022). This method achieves viroid multiplication at detectable titers by molecular techniques, confirming the results and providing greater specificity in diagnosis (Duran-Vila *et al.*, 1993). Therefore, the results obtained in this evaluation were confirmed using molecular methods, specifically RT-PCR, as a complementary diagnostic approach.

Molecular diagnosis

The RNA extraction performed using the protocol developed by Chang *et al.* (1993) yielded concentrations ranging from 564.7 to 2,698.42 ng μL^{-1} . The 260/280 absorbance ratios ranged from 1.99 to 2.23, and the 260/230 absorbance ratios varied between 1.92 and 2.41 (Fig. 3). These values are considered within the optimal ranges for use in laboratory practices, according to guidelines established by Koetsier and Cantor (2019).




Figure 3. Box plots of quantification and quality of RNA extracted from Etrog citron using the protocol of Chang *et al.* (1993), measured by spectrophotometry. A) concentration. B) purity ratios A260/280 and A260/230.

Amplification of the 18S ribosomal RNA (rRNA) region produced a fragment of approximately 250 base pairs (bp) in all analyzed samples (Fig. 4A). This result is consistent with the size reported by Du *et al.* (2006), indicating a synthesis of high-quality cDNA. In positive control (healthy Etrog citron plants inoculated with an HSVd isolate) RT-PCR allowed the amplification of a region of the HSVd viroid of approximately 174 bp which comprises the entire TL and P domains and partially the C domain, as reported by Bernad and Duran-Vila (2006). Non-specific amplified products exceeding the expected size were not included in the diagnosis. The viroid was not detected in either the 17 citrus varieties evaluated or the negative control (non-inoculated Etrog citron plant) (Fig. 4B). Thereby, the results obtained in the molecular diagnosis confirmed the healthiness in the 17 citrus varieties, consistent with the results obtained in the biological diagnosis.




Figure 4. Detection of 18S rRNA and HSVd by RT-PCR. A) Amplification of the 18S rRNA region. B) HSVd amplification by RT-PCR. M: Molecular weight marker 100 bp (ThermoScientific ref. #SM0323). (-): Negative control. (+): Positive control. (R): PCR reaction control without template. Lane 1: Minneola Tangelo, Line 2: Orlando Tangelo, Line 3: Clemenules mandarin, Line 4: ICA Bolo mandarin, Line 5: Sweety Orange, Line 6: Key lime, Line 7: Tahiti lime, Line 8: Pomelo. * Primer dimer.

RT-PCR is the most widely used method for diagnosing viroids in citrus, whether performed singly or in multiplex (Wang *et al.*, 2009; Camps *et al.*, 2014; Lin *et al.*, 2015; Rodríguez-Mora *et al.*, 2015). This technique is fast, economical, and effective, contributing to the improved detection and characterization of viroids affecting citrus. It also allows for the discrimination between pathogenic and non-pathogenic variants of HSVd (Bernad and Duran-Vila, 2006). The use of RT-PCR extends to indexing procedures, serving as an essential methodology for the early detection of viroids in citrus within certification programs (NAPPO, 2013).

Detecting HSVd infection in citrus cultivars poses significant challenges since infected plants can be asymptomatic and go unnoticed. Therefore, the application of both biological and molecular diagnostic methods becomes crucial for reliably monitoring the healthiness of citrus planting material.

This study confirmed the absence of HSVd in the citrus propagation material within the foundation category, including tangelos, grapefruits, mandarins, oranges, and acid limes. These

were maintained under controlled conditions within an anti-aphid mesh greenhouse at the Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de Investigación Palmira. Previous studies by Rodríguez-Mora *et al.* (2017) reported that this collection is free from systemic diseases such as tristeza, exocortis, and citrus huanglongbing, all regulated in the Colombian citrus certification standard (ICA, 2019). This achievement represents a significant milestone and provides substantial support to Colombia's citrus certification program.

CONCLUSION

This study confirmed the healthiness of the citrus variety collection within the foundation category regarding HSVd. This implies the potential use of this material as a reliable source of buds for plant multiplication within the registered and certified seed category. Thus, it actively contributes to maintaining quality and health standards in citrus nurseries in Colombia. This represents a significant step towards ensuring the healthy and sustainable production of new citrus cultivated areas.

Acknowledgments

These results are part of the project "Development of indexing processes for planting material against major diseases to ensure citrus health in Colombia, ID 1001260," funded by the Ministerio de Agricultura y Desarrollo Rural (MADR) and executed by the Corporación Colombiana de Investigación Agropecuaria – Agrosavia, Centro de Investigación Palmira.

Conflict of interests: The manuscript was prepared and reviewed with the participation of the authors, who declare that there is no conflict of interest that jeopardizes the validity of the presented results.

Author's contributions: D. M. Rodríguez-Mora: Methodological design and establishment of greenhouse and laboratory assays, writing the original draft, K. A. Rodríguez-Arévalo: establishment of greenhouse and laboratory assays, M. Martínez: Data analysis, and N. Murcia-Riaño: Funding acquisition, activities oversight, data analysis, manuscript review and editing. All authors have reviewed the final version of the manuscript.

BIBLIOGRAPHIC REFERENCES

Barbosa, C.J., J.A. Pina, J. Pérez-Panadés, L. Bernad, P. Serra, L. Navarro, and N. Duran-Vila. 2005. Mechanical transmission of citrus viroids. *Plant Dis.* 89(7), 749-754. Doi: <https://doi.org/10.1094/PD-89-0749>

Belabess, Z., N. Radouane, T. Sagouti, A. Tahiri, and R. Lahlali. 2021. A current overview of two viroids prevailing in citrus orchards: Citrus exocortis viroid and Hop stunt viroid. pp. 589-694. In: Khan, M.S. and I.A. Khan (eds.). *Citrus - research, development and biotechnology*. IntechOpen, London. Doi: <https://doi.org/10.5772/intechopen.95914>

Bernad, L. and N. Duran-Vila. 2006. A novel RT-PCR approach for detection and characterization of citrus viroids. *Mol. Cell. Probes* 20(2), 105-113. Doi: <https://doi.org/10.1016/j.mcp.2005.11.001>

Bové, J.M. 2006. Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. *J. Plant Pathol.* 88(1), 7-37.

Camps, R., M. Castro, and X. Besoain. 2014. Simultaneous detection of CTV, CEVd and HSVd using Arizona 861 S1 Citron and RT-PCR. *Cien. Inv. Agr.* 41(2), 255-261. Doi: <https://doi.org/10.4067/S0718-16202014000200012>

Chang, S., J. Puryear, and J. Cairney. 1993. A simple and efficient method for isolating RNA from pine trees. *Plant Mol. Biol. Rep.* 11(2), 113-116. Doi: <https://doi.org/10.1007/BF02670468>

Dawson, W.O., M. Bar-Joseph, S.M. Garnsey, and P. Moreno. 2015. Citrus tristeza virus: making an ally from an enemy. *Annu. Rev. Phytopathol.* 53, 117-155. Doi: <https://doi.org/10.1146/annurev-phyto-080614-120012>

Du, Z., J. Chen, and C. Hiruki. 2006. Optimization and application of a multiplex RT-PCR system for simultaneous detection of five potato viruses using 18S rRNA as an internal control. *Plant Dis.* 90(2), 185-189. Doi: <https://doi.org/10.1094/PD-90-0185>

Duran-Vila, N. 1989. Enfermedades producidas por viroides: la exocortis de los cítricos. *Phytoma* (7), 19-25.

Duran-Vila, N. 2004. Enfermedades de cítricos causadas por viroides: exocortis y caquexia. *Vida Rural* (188), 52-56.

Duran-Vila, N., J.A. Pina, J.F. Ballester, J. Juárez, C.N. Roistacher, R. Rivera-Bustamante, and J.S. Semancik. 1988. The citrus exocortis disease: a complex of viroid-RNAs. pp. 152-164. Proc 10th Conf. Int. Org. Citrus Virol. IOCV, Riverside CA. Doi: <https://doi.org/10.5070/C54CF7N8QB>

Duran-Vila, N., J.A. Pina, and L. Navarro. 1993. Improved indexing of citrus viroids. pp. 202-211. In: Moreno P., J.V. da Graca, and L.W. Timmer (eds.) Proc. 12th Conf. Int. Organ. Citrus Virol. IOCV. Riverside, CA. Doi: <https://doi.org/10.1023/A:1008711402856>

Duran-Vila, N., J.S. Semancik, and P. Broadbent. 2000. Viroid diseases, cachexia and exocortis. pp. 51-54. In: Timmer, L.W., S.M. Garnsey, and J.H. Graham (eds.). Compendium of citrus diseases. 2nd ed. APS, St. Paul, MN.

Eiras, M., S.R. Silva, E.S. Stuchi, S.A. Carvalho, and R.M. Garcêz. 2013. Identification and characterization of viroids in 'Navelina ISA 315' sweet orange. *Trop. Pant Pathol.* 38(1), 58-62. Doi: <https://doi.org/10.1590/S1982-56762013000100009>

Guerrero, C.E., O.G. Alvarado, H. Gutiérrez, R. Gozález, M.G. Álvarez, and M. Luna. 2013. Detección por RT-PCR punto final y tiempo real de tres especies de viroides en cítricos de Nuevo León y Tamaulipas, México. *Rev. Mex. Fitopatol.* 31(1), 20-28.

ICA, Instituto Colombiano Agropecuario. 2019. Resolución 12816, Por medio de la cual se establece los requisitos para el registro ante el ICA de los viveros y/o huertos básicos productores y/o comercializadores de semilla sexual y/o asexual (material vegetal de propagación) de cítricos,

así como los requisitos fitosanitarios para la conservación, producción, certificación y distribución de material de propagación de cítricos en viveros, en el territorio nacional. Bogota.

Koetsier, G. and E. Cantor. 2019. A practical guide to analyzing nucleic acid concentration and purity with microvolume spectrophotometers. N. Engl. Biolabs 1, 1-8.

Lin, C.Y., M.-L. Wu, T.-L. Shen, H.-H. Yeh, and T.-H. Hung. 2015. Multiplex detection, distribution, and genetic diversity of *Hop stunt viroid* and *Citrus exocortis viroid* infecting citrus in Taiwan. Virol. J. 12, 11. Doi: <https://doi.org/10.1186/s12985-015-0247-y>

MADR, Ministerio de Agricultura y Desarrollo Rural. 2021. Cadena de los cítricos. Indicadores e instrumentos. Segundo trimestre 2021. In: Sistema de información de gestión y desempeño de organizaciones de cadenas – SIOC, <https://sioc.minagricultura.gov.co/Citricos/Documentos/2021-06-30%20Cifras%20Sectoriales.pdf> ; consulted: May, 2023.

Mosquera, V., M.F. Martínez, W.J. Cuellar, J. Vaca-Vaca, I. Lozano, and N. Murcia. 2015. Detection of viroids and Citrus tristeza virus (CTV) in Tahití Lime *Citrus latifolia* (Tanaka) Through application of RT-PCR. p. 59. In: Libro de resúmenes, XXXIII Congreso Colombiano de Fitopatología y ciencias afines. Palmira, Colombia.

Murcia, N., L. Bernad, A. Caicedo, and N. Duran-Vila. 2010. Citrus viroids in Colombia. pp. 158-166. In: Proc. 17th Int. Org. Citrus Virol. IOCV. Riverside, CA. Doi: <https://doi.org/10.5070/C5008577NK>

Murcia, N., M. Betancourt, L. Pérez, D.M. Rodríguez, L. Ríos, Y.C. Pisco, and M.F. Martínez. 2020. Principales enfermedades en el cultivo de lima ácida Tahití. pp. 257-325. In: Castillo-Montaña, P. and L. Gaona. (ed.). Modelo productivo de lima ácida Tahití (*Citrus × latifolia* Tanaka ex Q. Jiménez) para Colombia. Agrosavia, Mosquera, Colombia. Doi: <https://doi.org/10.21930/agrosavia.model.7403435>

NAPPO, North American Plant Protection Organization. 2013. NRMF 16. Medidas integradas para la movilización de material propagativo de cítricos. Ottawa.

Ohno, T., N. Takamatsu, T. Meshi, and Y. Okada. 1983. Hop stunt viroid: molecular cloning and nucleotide sequence of the complete cDNA copy. Nucleic Acid Res. 11(18), 6184-6196. Doi: <https://doi.org/10.1093/nar/11.18.6185>

Rakowski, A.G., J.A. Szychowski, Z.S. Avena, and J.S. Semancik. 1994. Nucleotide sequence and structural features of the group III citrus viroid. J. Gen. Virol. 75(12), 3581-3584. Doi: <https://doi.org/10.1099/0022-1317-75-12-3581>

Reanwarakorn, K. and J.S. Semancik. 1998. Regulation of pathogenicity in hop stunt viroid related group II citrus viroids. J. Gen. Virol. 79(12), 3163-3171. Doi: <https://doi.org/10.1099/0022-1317-79-12-3163>

Reanwarakorn, K. and J.S. Semancik. 1999. Correlation of Hop stunt viroid variants to cachexia and xyloporosis disease of citrus. Phytopathol. 89(7), 568-574. Doi: <https://doi.org/10.1094/PHYTO.1999.89.7.568>

Rodríguez-Mora, D., V. Mosquera, M. Martínez, and N. Murcia. 2015. RT-PCR multiplex for detection of viroids in Tahiti acid lime (*Citrus latifolia* Tanaka). p 60. In: Libro de resúmenes, XXXII Cong. Colomb. Fitopatol. and I Simp. Int. Fusarium. Cajica, Colombia.

Rodríguez-Mora, D., L. Palacios, M. Martínez, and N. Murcia. 2017. Collection of work of citrus varieties free of tristeza, exocortis, and huanglongbing. p 35. En: Libro de resúmenes, V Simp. Int. Frutic. Trop. Subtropical and IX Simp. Int. Piña “Fruticultura”. La Habana.

Rodríguez-Mora, D., L. Palacios, M.F. Martínez, and N. Murcia. 2022. Manejo de enfermedades de cítricos en ambiente protegido. pp. 151-216. In: Gaona-García, L. (ed.). Bases tecnológicas para la producción de material de siembra de alta calidad de las especies cítricas

para Colombia. Agrosavia, Mosquera, Colombia. Doi:
<https://doi.org/10.21930/agrosavia.manual.7405682>

Semancik, J.S., C.N. Roistacher, R. Rivera-Bustamante, and N. Duran-Vila. 1988. Citrus cachexia viroid, a new viroid of citrus: relationship to viroids of the exocortis disease complex. J. Gen. Virol. 69(12), 3059-3068. Doi: <https://doi.org/10.1099/0022-1317-69-12-3059>

Semancik, J.S. and L.G. Weathers. 1972. Exocortis disease: evidence for a new species of “infectious” low molecular weight RNA in plants. Nature New Biol. 237, 242-244. Doi: <https://doi.org/10.1038/newbio237242a0>

Serra, P., S. Gago, and N. Duran-Vila. 2008. A single nucleotide change in Hop stunt viroid modulates citrus cachexia symptoms. Virus Res. 138(1-2), 130-134. Doi: <https://doi.org/10.1016/j.virusres.2008.08.003>

Serra, P., A. Messmer, D. Sanderson, D. Sanderson, D. James, and R. Flores. 2018. Apple hammerhead viroid-like RNA is a *bona fide* viroid: autonomous replication and structural features support its inclusion as a new member in the genus *Pelamoviroid*. Virus Res. 249, 8-15. Doi: <https://doi.org/10.1016/j.virusres.2018.03.001>

Teymuri, P., S. Nasrollanejad, V. Alavi, and M. Taghinasab. 2022. Biological characteristics of *Citrus cachexia viroid* isolates in local tangerine orchards (*Citrus unshiu* Marcovitch) in East Mazandaran. Appl. Entomol. Phytopathol. 90(1), 87-97. Doi: <https://doi.org/10.22092/JAEP.2022.359220.1444>

Vamenani, R., H. Rahimian, S.M. Alavi, and V. Babaeizad. 2014. Detection of hop stunt viroid in lemon, sweet lime, clementine and satsuma mandarins and grapefruit trees in mazandaran. Iran. J. Plant Path. 50(2), 105.

Venkataraman, S., U. Badar, E. Shoeb, G. Hashim, M. AbouHaidar, and K. Hefferon. 2021. An inside look into biological miniatures: molecular mechanisms of viroids. Int. J. Mol. Sci. 22(6), 2795. Doi: <https://doi.org/10.3390/ijms22062795>

Vernière, C., X. Perrier, C. Dubois, A. Dubois, L. Botella, C. Chabrier, J.M. Bové, and N. Duran Vila. 2004. Citrus viroids: symptom expression and effect on vegetative growth and yield of clementine trees grafted on Trifoliolate orange. Plant Dis, 88(1), 1189-1197. Doi: <https://doi.org/10.1094/PDIS.2004.88.11.1189>

Villalobos, W., L. Moreira, V.H. Alfaro, G. Yglesias, and C. Rivera. 2004. Detección del Viroide de la exocortis en una plantación comercial de naranja dulce, utilizando la selección 861-S1 del cidro 'Etrog' (*Citrus medica* L.). Manejo Integrado de Plagas y Agroecología 72, 86-90.

Wang, X., Y. Zhou, Z. Li, K. Tang, Y. Liu, M. Cao, and C. Zhou. 2010. Molecular, biological and phylogenetic analysis of chinese isolates of *Hop stunt viroid* associated with citrus cachexia disease. J. Phytopathol. 158(5), 372-377. Doi: <https://doi.org/10.1111/j.1439-0434.2009.01629.x>

Wang, X., C. Zhou, Z. Tang, Y. Zhou, and Z. Li. 2009. A rapid one-step multiplex RT-PCR assay for the simultaneous detection of five citrus viroids in China. Eur. J. Plant Pathol. 124, 175-180. Doi: <https://doi.org/10.1007/s10658-008-9386-y>

Zeitoni, H., S.M.B. Hashemian, and M. Shams-Bakhsh. 2023. Detection of hop stunt viroid variants from naturally infected kumquat and limequat trees in Mazandaran Province, Iran. J. Phytopathol. 105, 545-556. Doi: <https://doi.org/10.1007/s42161-023-01347-8>