

Acaricidal and larvicidal activity of *Piper marginatum* and *Piper tuberculatum* essential oils from Ecuador

Actividad acaricida y larvicida de los aceites esenciales de *Piper marginatum* y *Piper tuberculatum* de Ecuador



SHIRLEY MADELEINE MONCAYO-BAÑO^{1, 2, 4}
MARÍA EUGENIA RONDÓN-RIVAS²
FRANCISCO XAVIER CORNEJO-SOTOMAYOR³

Larvae of *A. aegypti* and *R. microplus* after treatment with essential oils of *P. marginatum* (top) and *P. tuberculatum* (bottom).

Photo: S.M. Moncayo-Baño

ABSTRACT

Piper marginatum and *Piper tuberculatum* are species used for their insecticidal properties by local indigenous people in eastern Ecuador. This study aimed to determine the acaricidal and larvicidal activity of the essential oil of these species. The essential oils were obtained by hydrodistillation. The acaricidal activity was evaluated with the contact toxicity method against larvae of the tick *Rhipicephalus microplus*. Larvicidal activity was evaluated against *Aedes aegypti* mosquito larvae following WHO recommended guidelines. The essential oils of *P. marginatum* and *P. tuberculatum* showed an acaricidal potential with LC₅₀ of 0.90 and 0.73 $\mu\text{L mL}^{-1}$, respectively. In addition, excellent larvicidal activity was obtained with LC₅₀ of 11.87 $\mu\text{L mL}^{-1}$ (*P. marginatum*) and 8.42 $\mu\text{L mL}^{-1}$ (*P. tuberculatum*) at 48 hours of evaluation. This is the first report of the acaricidal activity of *P. marginatum* and *P. tuberculatum* essential oils against *R. microplus* tick larvae. The results indicate that essential oils could serve as a natural alternative for controlling these pests, with ethnobotanical applications as a lice and tick repellent.

Additional key words: *Piper*; larvae; *Rhipicephalus microplus*; *Aedes aegypti*; ethnobotanical.

¹ Universidad Agraria del Ecuador, Facultad de Ciencias Agrarias, Carrera Agroindustria, Guayaquil (Ecuador).
ORCID Moncayo-Baño, S.M.: <https://orcid.org/0000-0003-2374-497X>

² Universidad de Los Andes, Facultad de Farmacia y Bioanálisis, Cátedra de Farmacognosia, Merida (Venezuela).
ORCID Rondón-Rivas, M.E.: <https://orcid.org/0000-0003-2393-751X>

³ Universidad de Guayaquil, Facultad de Ciencias Naturales, Herbario GUAY, Guayaquil (Ecuador).
ORCID Cornejo-Sotomayor, F.X.: <https://orcid.org/0000-0002-4081-4047>


⁴ Corresponding author. smoncayo@uagraría.edu.ec

RESUMEN

Piper marginatum y *Piper tuberculatum* son especies utilizadas por sus propiedades insecticidas por indígenas locales del este de Ecuador. Este estudio tuvo como objetivo determinar la actividad acaricida y larvicida del aceite esencial de estas especies. Los aceites esenciales se obtuvieron por hidrodestilación. La actividad acaricida se evaluó con el método de toxicidad por contacto frente a larvas de la garrapata *Rhipicephalus microplus*. Se evaluó la actividad larvicida contra larvas del mosquito *Aedes aegypti* siguiendo los lineamientos recomendados por la OMS. Los aceites esenciales de *P. marginatum* y *P. tuberculatum* mostraron un potencial acaricida con CL_{50} de 0.90 y 0.73 $\mu\text{L mL}^{-1}$, respectivamente. Además, se obtuvo una excelente actividad larvicida con CL_{50} de 11.87 $\mu\text{L mL}^{-1}$ (*P. marginatum*) y 8.42 $\mu\text{L mL}^{-1}$ (*P. tuberculatum*) a las 48 horas de evaluación. Este es el primer reporte de la actividad acaricida de los aceites esenciales de *P. marginatum* y *P. tuberculatum* frente a las larvas de la garrapata *R. microplus*. Los resultados indican que los aceites esenciales podrían servir como una alternativa natural para el control de estas plagas, con aplicaciones etnobotánicas como repelente de piojos y garrapatas.

Palabras clave adicionales: *Piper*; larvas; *Rhipicephalus microplus*; *Aedes aegypti*; etnobotánica.

Received: 17-03-2024 Accepted: 28-04-2024 Published: 24-05-2024



INTRODUCTION

The most significant tick in livestock worldwide is *Rhipicephalus (Boophilus) microplus*. This hematophagous ectoparasite is one of the main obstacles to the productivity of the livestock industry in areas where this species is endemic (Nicaretta *et al.*, 2021). It causes a decrease in cattle weight gain, damage to hides, reduced meat and milk production, myiasis and transmission of zoonotic diseases (Polanco and Ríos, 2016). Due to its significance, it is the most studied species in acaricide trials (Adenubi *et al.*, 2018). *Aedes aegypti* are vectors of arboviral pathogens that annually cause hundreds of millions of cases of Zika, dengue fever, yellow fever, and chikungunya. They thrive in urbanized areas where they are in close contact with people, making them an exceptionally successful vector. *A. aegypti* is extremely common in areas that lack piped water systems and rely heavily on stored water for breeding sites (Matthews, 2019; Zettel and Kaufman, 2019).

In the prevention of vector-borne diseases, there is a trend to develop alternative methods based on natural products to replace chemical pesticides and delay the development of resistance, a problem that causes millions of dollars in losses (Selles *et al.*, 2021). Essential oils are metabolites formed by a complex mixture of several chemical compounds mainly terpenes and phenylpropanoids, which give plants their characteristic aroma and possess a wide range of biological properties, including

antimicrobial, antifungal and insecticidal activities (De Souza *et al.*, 2023). Studies conducted on the essential oils of *Piper* species (Piperaceae) highlight their great potential for pest control and the low environmental impact observed during the process (Pereira *et al.*, 2021). Their efficacy against different species of mosquitoes and mites at low concentrations has been reported (Huong *et al.*, 2019; Araújo *et al.*, 2020).

Piper marginatum Jacq. and *Piper tuberculatum* Jacq. are native neotropical species that stand out for their pharmacological potential and are widely used in folk medicine for their healing properties (Salehi *et al.*, 2019). *P. marginatum* is known in Ecuador as cordoncillo blanco, Santa María and "Anotede" in Tsa'fiki language (De la Torre, 2008). Its essential oils have shown antibacterial, antioxidant, antiparasitic and larvicidal activities (Ayres *et al.*, 2021). *P. tuberculatum* is used in Ecuador to combat pediculosis problems (De la Torre *et al.*, 2008). Previous studies have shown that its essential oils have antimicrobial, insecticidal and antiprotozoal effects (Da Silva *et al.*, 2023).

This study reports the acaricidal and larvicidal activity of the essential oils of *P. marginatum* and *P. tuberculatum* against larvae of the tick *R. microplus* and the mosquito *A. aegypti*. These species cause serious damage to human and animal health and result in serious economic losses.

MATERIALS AND METHODS

Plant materials

Leaves, stems, and inflorescences of *P. marginatum* and *P. tuberculatum* were collected in Cumanda, Chimbo-razo province (2°12'12" S, 79°06'36" W) and on the El Triunfo - Bucay road, Guayas province (2°18'9" S, 79°14'16" W) from Ecuador. The identification of the botanical species was carried out in the GUAY Herbarium of the Faculty of Natural Sciences of the University of Guayaquil, Ecuador, where the vouchers of the specimens were deposited under the codes MER06 (*P. marginatum*) and MER07 (*P. tuberculatum*).

Hydrodistillation

The leaves, stems, and inflorescences of the species were subjected to hydrodistillation for 3 h, using a Clevenger-type apparatus. The oils, once obtained, were meticulously dried over anhydrous sodium sulfate and then stored in sealed vials. These vials were then placed in a controlled environment at 4°C in the dark to ensure the oils' stability and integrity until they were ready for analysis and testing.

Acaricidal activity

Acaricidal activity was analyzed using the contact toxicity technique described by Tabari *et al.* (2020) with minor modifications. Larvae between 14 and 21 d old of the tick *R. microplus* obtained from gravid females collected from infested cattle were used. The tick species was identified by Prof. Antonio Ascensão of the Laboratory of Biodiversity of Arthropods, Department of Biology, Faculty of Sciences, University of Los Andes, Venezuela. Serials of essential oil dilutions were prepared from 20.0 to 0.3 $\mu\text{L mL}^{-1}$, in 2% Tween® 80 solution. A Tween® 80 solution was used as a negative control and a 20 $\mu\text{L mL}^{-1}$ cypermethrin solution dissolved in Tween® 80 (2%), prepared from the commercial product CIPER-VEEX® (10% cypermethrin), was used as a positive control. Whatman® N°1 filter paper discs were impregnated with 1 mL of the test solutions. Tick larvae were transferred to two of the impregnated paper disks, placed inside a Petri dish. The plates were sealed and stored in a glass chamber at 27±2°C and 85±2% humidity. Assays were performed in duplicate. Mortality records were made after 48 h under a stereomicroscope.

Larvicidal activity

Larvicidal activity was analyzed following the guidelines of the World Health Organization (WHO, 1981) for third and fourth instar larvae of wild *A. aegypti*, collected at the Faculty of Natural Sciences of the University of Guayaquil. The larvae were identified by the author using taxonomic guides. An aliquot of essential oils was solubilized in absolute ethanol until a stock solution was obtained, to prepare test solutions in serial dilutions of 3 to 1,000 $\mu\text{L mL}^{-1}$. The final volume of the solutions was 50 mL. A total of 10 larvae were exposed to the test solutions. The negative control consisted of a 2% ethanol solution; the positive control consisted of a solution of the organophosphate insecticide malathion (1 $\mu\text{L mL}^{-1}$). Each concentration was evaluated in triplicate. The assays were stored at 25±2°C and relative humidity of 70±2%. Larval mortality readings were taken at 24 and 48 h.

Statistical analyses

The concentrations producing 50 and 95% mortality in the mite and larvae population (LC_{50}) and (LC_{95}) and their respective confidence limits (95%) were determined by Probit analysis. To establish statistically significant differences between the assays, a one-way ANOVA analysis of variance was performed using the Fisher (LSD) method. Values of $P < 0.05$ were considered significant. All the statistical analyses were carried out using Statgraphics Centurion XVI statistical software.

RESULTS AND DISCUSSION

Acaricidal activity

The mortality percentages of *R. microplus* larvae after 48 h of exposure to the essential oils of *P. marginatum* and *P. tuberculatum* are shown in tables 1 and 2. No mortality was reported in the negative control, while mortality with the positive control was 100%. The results indicate that the essential oil of *P. tuberculatum* was the most toxic, causing 100% mortality of *R. microplus* larvae at a concentration of 1 $\mu\text{L mL}^{-1}$. The ANOVA comparison analysis found statistically significant differences between the lethal concentrations of the essential oils at 95% significance level.

Table 1. Mortality percentage of *Rhipicephalus microplus* caused by different concentrations of *Piper marginatum* essential oil.

Concentration ($\mu\text{L mL}^{-1}$)	0.6	1.2	2.5	5.0	10.0	20.0
Mortality	11 \pm 2	86 \pm 1	92 \pm 3	100 \pm 0	100 \pm 0	100 \pm 0

Mean of $n=3\pm$ standard deviation.

Table 2. Mortality percentage of larvae of *Rhipicephalus microplus* caused by different concentrations of *Piper tuberculatum* essential oil.

Concentration ($\mu\text{L mL}^{-1}$)	0.3	0.6	0.8	1.0	2.5
Mortality	6 \pm 1	23 \pm 2	49 \pm 1	100 \pm 0	100 \pm 0

Mean of $n=3\pm$ standard deviation.

Table 3 summarizes the LC_{50} and LC_{95} values of *P. marginatum* and *P. tuberculatum* oils against *R. microplus* tick larvae. The most potent oil was that of *P. tuberculatum*, with a LC_{50} of $0.73 \mu\text{L mL}^{-1}$ while the oil of *P. marginatum* showed an LC_{50} of $0.90 \mu\text{L mL}^{-1}$.

Table 3. Lethal concentrations of the essential oils of *Piper* species on larvae of *Rhipicephalus microplus*.

Essential oil	LC_{50}	LC_{95}
<i>P. marginatum</i>	0.90 (0.75-1.02)	2.10 (1.86-2.47)
<i>P. tuberculatum</i>	0.73 (0.71-0.75)	1.06 (1.02-1.11)

(95% CI) = 95% confidence interval

This groundbreaking report unveils the acaricidal activity of essential oils from *Piper* species against *R. microplus* larvae, a discovery of utmost significance as larvae are pivotal in tick control systems. Da Silva *et al.* (2017) suggest that essential oils from plants of the genus *Piper*, rich in phenylpropanoids and hydrocarbonated sesquiterpenes, as in this study, exhibit potent insecticidal and acaricidal activity.

The practical implications of this research are evident in the evaluation of the essential oil of *P. marginatum* and *P. tuberculatum* against the mite *Tetranychus urticae*, a crop pest from the family Tetranychidae. The obtained LC_{50} values of 0.90 and $0.50 \mu\text{L mL}^{-1}$, respectively (Ribeiro *et al.*, 2016; Araújo *et al.*, 2020) underscore the potential of these essential oils in pest control.

According to Selles *et al.* (2021), the acaricidal activity of essential oils is related to their hydrophobic nature, which is responsible for several mechanical effects, causing death by water stress or asphyxia due

to alteration of cuticular waxes and blockage of respiratory stigmas. Essential oils have been shown to act on the motor function of *R. microplus* larvae, increasing catecholamine levels within the central nervous system (Salman *et al.*, 2020) and inhibiting the action of acetylcholinesterase in this tick species (Cardoso *et al.*, 2020).

The acaricidal capacity observed in the extracts supports the ethnobotanical use of *P. tuberculatum* in Ecuador as a lice and tick repellent (De la Torre *et al.*, 2008).

Larvicidal activity

The concentrations of the essential oils and their respective percentages of mortality after 24 and 48 h of exposure are shown in tables 4 and 5. *A. aegypti* larvae's mortality percentage was 100% from the concentration of $50 \mu\text{L mL}^{-1}$ in both oils evaluated. No mortality was reported in the negative control, while mortality with the positive control was 100%.

The lethal concentrations of the essential oils are shown in table 6. *P. tuberculatum* obtained the highest activity with LC_{50} of $8.42 \mu\text{L mL}^{-1}$ and LC_{95} of $51.38 \mu\text{L mL}^{-1}$ while *P. marginatum* obtained LC_{50} of $11.87 \mu\text{L mL}^{-1}$ and LC_{95} of $28.29 \mu\text{L mL}^{-1}$ at 48 h of exposure. The ANOVA comparison analysis found statistically significant differences between the lethal concentrations of the essential oils at 95% significance level.

The volatile compounds of *Piper* species are characterized by their great potential for pest control and the low environmental impact (Pereira *et al.*, 2021). Previous larvicidal evaluations on the oils of *P. marginatum*

Table 4. Mortality percentage of *Aedes aegypti* larvae caused by different concentrations of *Piper marginatum* essential oil ($\mu\text{L mL}^{-1}$).

Exposure time (h)	3	6	12	16	20	25	50-250
24	10 \pm 0	27 \pm 6	30 \pm 0	53 \pm 6	73 \pm 6	83 \pm 6	100 \pm 0
48	17 \pm 6	30 \pm 6	43 \pm 6	67 \pm 6	87 \pm 15	90 \pm 0	100 \pm 0

Mean of $n=3 \pm$ standard deviation.

Table 5. Mortality percentage of *Aedes aegypti* larvae caused by different concentrations of *Piper tuberculatum* essential oil ($\mu\text{L mL}^{-1}$).

Exposure time (h)	3	6	12	25	30	40-250
24	17 \pm 6	30 \pm 10	33 \pm 6	60 \pm 0	70 \pm 10	100 \pm 0
48	30 \pm 10	57 \pm 6	60 \pm 10	70 \pm 0	80 \pm 10	100 \pm 0

Mean of $n=3 \pm$ standard deviation.

Table 6. Lethal concentrations (LC) of the essential oils of *Piper marginatum* and *P. tuberculatum* species on 3^o- 4^o instar larvae of the mosquito *Aedes aegypti*.

Exposure time (h)	LC	<i>P. marginatum</i>	<i>P. tuberculatum</i>
24	LC ₅₀ (CI)	14.65 (11.90-17.84)	19.70 (15.43-25.53)
	LC ₉₅ (CI)	32.34 (26.95-42.53)	52.69 (41.40-78.73)
48	LC ₅₀ (CI)	11.87 (9.21-14.61)	8.42 (4.25-13.08)
	LC ₉₅ (CI)	28.29 (23.63-36.83)	51.38 (37.76-94.06)

(95% CI) = 95% confidence interval.

and *P. tuberculatum* (Lavor *et al.*, 2012; Pereira *et al.*, 2021), reported LC₅₀ of 39 and 106 $\mu\text{L mL}^{-1}$, respectively. Essential oils from Ecuadorian species showed higher larvicidal activity, probably due to their different chemical composition and synergistic interactions between their compounds. The plant extracts used in the present study were previously characterized to identify their phytochemical composition by our research team. Compounds such as sesquiterpenes curzerene (22.44%) and atractylone (18.15%) were identified in *P. marginatum*, while in *P. tuberculatum*, the phenylpropanoid dilapiol (49.15%) was identified (Moncayo *et al.*, 2021). Species with essential oils predominantly composed of these compounds have shown insecticidal properties against both adult and larvae stages of *A. aegypti* and *Culex quinquefasciatus* mosquitoes (Benelli *et al.*, 2017; Hung *et al.*, 2020; Pereira *et al.*, 2021). Dilapiol, a potent inhibitor of cytochrome activity (P450 and CYP34A), reduces the ability to excrete xenobiotics causing the insect's death by accumulation of toxic substances in the

digestive tract (Santos *et al.*, 2021). In addition, its structure can amplify the effects of other chemicals (Durofil *et al.*, 2021).

CONCLUSION

The present study demonstrated that the essential oils of *P. marginatum* and *P. tuberculatum* possess high acaricidal (*R. microplus*) and larvicidal (*A. aegypti*) activity and could serve as a natural alternative for the biocontrol of these dangerous disease vectors. This is the first report on the activity of *P. marginatum* and *P. tuberculatum* essential oils against *R. microplus* tick larvae.

Conflict of interests: The manuscript was prepared and reviewed with the participation of the authors, who declare that there is no conflict of interest that puts at risk the validity of the presented results.

BIBLIOGRAPHIC REFERENCES

- Adenubi, O., A. Ahmed, F. Fasina, L. McGaw, J. Eloff, and V. Naidoo. 2018. Pesticidal plants as a possible alternative to synthetic acaricides in tick control: a systematic review and meta-analysis. *Ind. Crops Prod.* 123, 779-806. Doi: <https://doi.org/10.1016/j.indcrop.2018.06.075>
- Araújo, M., M. Moraes, F. Born, and C. Camara. 2020. Acaricidal property and repellent action against *Tetranychus urticae* Koch of essential oils from three species of *Piper* that occur in fragments of the Atlantic Forest in the State of Pernambuco, Brazil. *Rev. Virtual Quim.* 12(3), 629-638. Doi: <https://doi.org/10.21577/1984-6835.20200050>
- Ayres, V., M. Oliveira, E. Baldin, G. Corrêa, A. Guimarães, and R. Takeara. 2021. Chemical composition and insecticidal activity of the essential oils of *Piper marginatum*, *Piper callosum* and *Vitex agnus-castus*. *An. Acad. Bras. Cienc.* 93(3), 1-11. Doi: <https://doi.org/10.1590/0001-3765202120200616>
- Benelli, G., R. Pavela, R. Iannarelli, R. Petrelli, L. Cappellacci, K. Cianfaglione, F. Afshar, M. Nicoletti, A. Canale, and F. Maggi. 2017. Synergized mixtures of Apiaceae essential oils and related plant-borne compounds: larvicidal effectiveness on the filariasis vector *Culex quinquefasciatus* Say. *Ind. Crops Prod.* 96, 186-195. Doi: <https://doi.org/10.1016/j.indcrop.2016.11.059>
- Cardoso, A., E. Santos, A. Lima, K. Temeyer, A. Pérez de León, L. Costa, A. Junior, and A. Soares. 2020. Terpenes on *Rhipicephalus (Boophilus) microplus*: Acaricidal activity and acetylcholinesterase inhibition. *Vet. Parasitol.* 280, 109090. Doi: <https://doi.org/10.1016/j.vetpar.2020.109090>
- Da Silva, L., C. Paulo, T. Moura, D. Alves, R. Pessoa, I. Araújo, C. Morais, S. Tintino, C. Nonato, J. Costa, J. Ribeiro, H. Coutinho, G. Kowalska, P. Mitura, M. Bar, R. Kowalski, and I. Menezes. 2023. Antibacterial activity of the essential oil of *Piper tuberculatum* Jacq. fruits against multidrug-resistant strains: inhibition of efflux pumps and β -Lactamase. *Plants* 12(12), 2377. Doi: <https://doi.org/10.3390/plants12122377>
- Da Silva, J., R. Trindade, N. Alves, P. Figueiredo, J. Maia, and W. Setzer. 2017. Essential oils from neotropical *Piper* species and their biological activities. *Int. J. Mol. Sci.* 18(12), 2571. Doi: <https://doi.org/10.3390/ijms18122571>
- De la Torre, L., H. Navarrete, P. Muriel, M. Macía, and H. Balslev. 2008. Formato del catálogo de especies de plantas útiles del Ecuador. pp. 496-504. In: De la Torre, L., H. Navarrete, P. Muriel, M. Macía, and H. Balslev (eds.). *Enciclopedia de las plantas útiles del Ecuador*. Herbario QCA; Herbario AAU, Quito; Aarhus.
- De Souza, B., O. Oliveira, S. Mali, A. Anand, J. Nevez, C. Pereira, S. Kumar, R. Kumar, M. Moraes, M. Santana, and E. Aguiar. 2023. Chemical composition, preliminary toxicity, and antioxidant potential of *Piper marginatum sensu lato* essential oils and molecular modeling study. *Molecules* 28(15), 1-19. Doi: <https://doi.org/10.3390/moléculas28155814>
- Durofil, A., M. Radice, J. Blanco, and T. Ruiz-Téllez. 2021. *Piper aduncum* essential oil: a promising insecticide, acaricide and antiparasitic. A review. *Parasite* 28(42), 1-19.
- Hung, N., L. Huong, N. Chung, N. Thuong, P. Satyal, N. Dung, T. Tai, and W. Setzer. 2020. *Callicarpa* species from central Vietnam: essential oil compositions and mosquito larvicidal activities. *Plants* 9(113), 1-32. Doi: <https://doi.org/10.3390/plants9010113>
- Huong, L., N. Hung, D. Dai, T. Tai, V. Hien, P. Satyal, and W. Setzer. 2019. Chemical compositions and mosquito larvicidal activities of essential oils from *Piper* species growing wild in central Vietnam. *Molecules* 24(21), 1-30. Doi: <https://doi.org/10.3390/molecules24213871>
- Lavor, P., G. Santiago, R. Gois, L. Sousa, G. Bezerra, A. Arriaga, T. Lemos, P. Alvares, and P. Gomes. 2012. Larvicidal activity against *Aedes aegypti* of essential oils from northeast Brazil. *Nat. Prod. Commun.* 7(10), 1391-1392.
- Matthews, B.J. 2019. *Aedes aegypti*. *Trends Genet.* 35(6), 470-471. Doi: <https://doi.org/10.1016/j.tig.2019.03.005>
- Moncayo, S., M. Rondón, L. Araujo, L. Rojas, X. Cornejo, W. Guamán, and S. Jaramillo. 2021. Composición química y actividad biológica de los aceites esenciales de *Piper marginatum* Jacq. y *Piper tuberculatum* Jacq. de Ecuador. *Rev. Fac. Farm.* 63(1), 14-24.
- Nicaretta, J., D. Zapa, L. Couto, L. Heller, A. Cavalcante, L. Cruvinel, and W. Lopes. 2021. *Rhipicephalus microplus* seasonal dynamic in a Cerrado biome, Brazil: an update data considering the global warming. *Vet. Parasitol.* 296(109506), 1-10. Doi: <https://doi.org/10.1016/j.vetpar.2021.109506>
- Pereira, A., G. Pessoa, L. Yamaguchi, M. Stanton, A. Serravite, R. Pereira, W. Neves, and M. Kato. 2021. Larvicidal activity of essential oils from *Piper* species against strains of *Aedes aegypti* (Diptera: Culicidae) resistant to pyrethroids. *Front. Plant Sci.* 12(685864), 1-14. Doi: <https://doi.org/10.3389/fpls.2021.685864>
- Polanco, D. and L. Ríos. 2016. Aspectos biológicos y ecológicos de las garrapatas duras. *Cienc. Tecnol.* 17(1), 81-95. Doi: https://doi.org/10.21930/rcta.vol17_num1_art:463
- Ribeiro, N., C. Camara, and C. Ramos. 2016. Toxicity of essential oils of *Piper marginatum* Jacq. Against *Tetranychus urticae* Koch and *Neoseiulus californicus* (McGregor). *Chil. J. Agric. Res.* 76(1), 71-76.
- Salehi, B., Z. Zakaria, R. Gyawali, S. Ibrahim, J. Rajkovic, Z. Shinwari, T. Khan, J. Sharifi, A. Ozleyen, E. Turkdonmez, M. Valussi, T. Boyunegmez, L. Monzote, M. Martorell, and W. Setzer. 2019. *Piper*

- Species: A comprehensive review on their phytochemistry, biological activities and applications. *Molecules* 24(7), 1-118. Doi: <https://doi.org/10.3390/molecules24071364>
- Salman, M., R. Abbas, M. Israr, A. Abbas, M. Mehmood, M. Khan, Z. Sindhu, R. Hussain, M. Saleemi, and S. Shah. 2020. Repellent and acaricidal activity of essential oils and their components against *Rhipicephalus* ticks in cattle. *Vet. Parasitol.* 283(109178), 1-37.
- Santos, T., T. Vieira, J. Paula, J. Oliveira Neto, L. Cunha, L. A. Santos, and C. Romano. 2021. Influence of drying on the chemical composition and bioactivity of *Piper aduncum* (Piperaceae) essential oil against *Aedes aegypti* (Diptera: Culicidae). *Res. Soc. Dev.* 10(8), 1-10. Doi: <https://doi.org/10.33448/rsd-v10i8.17397>
- Selles, S., M. Koudri, M. González, J. González, M. Sánchez, A. González, J. Sanchis, L. Elhachimi, A. Olmeda, J. Tercero, and F. Valcárcel. 2021. Acaricidal and repellent effects of essential oils against ticks: a review. *Pathogens* 10(1379), 1-17. Doi: <https://doi.org/10.3390/pathogens10111379>
- Tabari, M., A. Rostami, A. Khodashenas, F. Maggi, R. Petrelli, C. Giordani, L. Taponjof, F. Papag, Y. Zuog, K. Cianfaglione, and M. Youssefi. 2020. Acaricidal activity, mode of action, and persistent efficacy of selected essential oils on the poultry red mite (*Dermanyssus gallinae*). *Food Chem. Toxicol.* 138(111207), 1-7. Doi: <https://doi.org/10.1016/j.fct.2020.111207>
- WHO, World Health Organization. 1981. Instructions for determining the susceptibility or resistance of mosquito larvae to insecticides. In: Division of Vector Biology and Control, <https://apps.who.int/iris/handle/10665/69615>; consulted: october, 2023.
- Zettel, C. and P. Kaufman. 2019. *Aedes aegypti*. In: Featured Creatures University of Florida, http://entnemdept.ufl.edu/creatures/aquatic/aedes_aegypti.htm#top; consulted: october, 2023.